

Mixing Colour(ing)s in Graphs

JAN VAN DEN HEUVEL

Centre for Discrete and Applicable Mathematics
Department of Mathematics
London School of Economics and Political Science



Wish we were there . . .





or there ...

- reporting on research by: **PAUL BONSMMA** (Twente)
LUIS CERECEDA (LSE)
JVDH (LSE)
and **MATTHEW JOHNSON** (Durham)

- in several different combinations

- and it all started with a question of
HAJO BROERSMA (Durham)

First definitions


- graph $G = (V, E)$: finite, simple, no loops, n vertices
- k -colouring of G : proper vertex-colouring
using colours from $\{1, 2, \dots, k\}$
 - we always assume $k \geq \chi(G)$
 - we use α, β, \dots to indicate k -colourings
- k -colour graph $\mathcal{C}(G; k)$
 - vertices are the k -colourings of G
 - two k -colourings are adjacent
if they differ in the colour on exactly one vertex of G

Some examples

- 2-colour graph for K_2 :



- 3-colour graph for K_2 :



Central question

General question

- Given G and k , what can we say about the colour graph $\mathcal{C}(G; k)$?

In particular

- is $\mathcal{C}(G; k)$ connected?

good way to think about it:

- can we go from any k -colouring to any other k -colouring by recolouring one vertex at the time?

Terminology: $\mathcal{C}(G; k)$ is **connected** $\iff G$ is k -mixing

Research on $\mathcal{C}(G; k)$

- little research in pure graph theory
- related to work in theoretical physics on Glauber dynamics of k -state anti-ferromagnetic Potts models at zero temperature
- related to work in theoretical computer science on Markov chain Monte Carlo methods for generating random k -colourings (“rapid mixing”)

Some first results on mixing

■ $k \geq \Delta(G) + 2 \implies G$ is k -mixing (‘‘Well-known’’)

■ $\underline{D(G)} = \max \{ \delta(H) \mid H \subseteq G \}$ (degeneracy)

Property


■ $k \geq D(G) + 2 \implies G$ is k -mixing (Dyer, et al., 2004)

Theorem

■ $k \geq D(G) + 3 \implies \mathcal{C}(G; k)$ is hamiltonian
(Choo & MacGillivray, 2006)

Extremal graphs for the degree bounds

- “boring” extremal graph: complete graph K_m
 - $\Delta(K_m) + 1 = D(K_m) + 1 = m$
 - all m -colourings look the same:
 - no vertex can change colour




Terminology

- frozen k -colouring: colouring in which no vertex can change colour
 - frozen colourings form **isolated vertices** in $\mathcal{C}(G; k)$
 - immediately mean G is not k -mixing

The case for planar graphs

■ G planar $\implies D(T) \leq 5 \implies k$ -mixing for $k \geq 7$

■ best possible :




frozen 6-colouring of the icosahedron

More interesting extremal graphs

- graph L_m : $K_{m,m}$ minus a perfect matching ($m \geq 3$)

- $\Delta(L_m) + 1 = D(L_m) + 1 = m$




- has frozen m -colourings – hence L_m is not m -mixing
- so:
bipartite graphs can be non- k -mixing for arbitrarily large k

More interesting properties of L_m

- non- k -mixing for $k = m$ colours
- but k -mixing for $3 \leq k \leq m - 1$
 - suppose L_m coloured with $k \leq m - 1$ colours

More interesting properties of L_m


- non- k -mixing for $k = m$ colours
- but k -mixing for $3 \leq k \leq m - 1$
 - suppose L_m coloured with $k \leq m - 1$ colours



- some colour c must appear more than once on the top

More interesting properties of L_m


- non- k -mixing for $k = m$ colours
- but k -mixing for $3 \leq k \leq m - 1$
 - suppose L_m coloured with $k \leq m - 1$ colours



- that colour c can't appear among the bottom vertices

More interesting properties of L_m


- non- k -mixing for $k = m$ colours
- but k -mixing for $3 \leq k \leq m - 1$
 - suppose L_m coloured with $k \leq m - 1$ colours



- so all vertices on the top can be recoloured to C

More interesting properties of L_m


- non- k -mixing for $k = m$ colours
- but k -mixing for $3 \leq k \leq m - 1$
 - suppose L_m coloured with $k \leq m - 1$ colours



- then the bottom can be recoloured to some $c' \neq c$

More interesting properties of L_m


- non- k -mixing for $k = m$ colours
- but k -mixing for $3 \leq k \leq m - 1$
 - suppose L_m coloured with $k \leq m - 1$ colours



- hence any colouring is connected to a 2-colouring
- easy to see that all these 2-colourings are connected

More interesting properties of L_m


- non- k -mixing for $k = m$ colours
- but k -mixing for $3 \leq k \leq m - 1$
 - suppose L_m coloured with $k \leq m - 1$ colours




- hence any colouring is connected to a 2-colouring
- easy to see that all these 2-colourings are connected
- so: mixing is not a monotone property

Mixing for small values of k

- smallest possible is $k = \chi(G)$
- $\chi(G) = 1$: graph without edges – boring
- $\chi(G) = 2$: bipartite graph with at least one edge
 - not-mixing for $k = 2$:



can't become



- $\chi(G) = 3$: 3-colourable graph with at least one odd cycle

The case $k = \chi = 3$



- cycle C_3 has six 3-colourings, all frozen
 $\implies C_3$ is not 3-mixing
- cycle C_5 has 30 3-colourings, none of them frozen
 - the colour graph $\mathcal{C}(C_5; 3)$ is formed of two 15-cycles
 $\implies C_5$ is not 3-mixing

Theorem

- $\chi(G) = 3 \implies G$ is not 3-mixing

Proof looks at 3-colourings of cycles


- suppose α is a 3-colouring of G
and C is a cycle in G
 - choose an **orientation** \vec{C} of the cycle
 - weight of an arc of \vec{C} :



- weight of the oriented cycle:
 $w(\vec{C}; \alpha) =$ sum of the weights of the arcs

Proof looks at 3-colourings of cycles

■ Example :



$$w(\vec{C}; \alpha) = -3$$

Weights of 3-colourings of cycles

- recolour one vertex to go from α to β



$$\implies w(\vec{C}; \alpha) = w(\vec{C}; \beta)$$

Property

- α and β connected by a path in $\mathcal{C}(G; 3)$

$$\implies \text{for all cycles } C \text{ in } G : w(\vec{C}; \alpha) = w(\vec{C}; \beta)$$

Weights of 3-colourings of cycles

- given 3-colouring α , form α^* by swapping colours 1 and 2

\implies all arcs change sign

\implies so for all C in G : $w(\vec{C}; \alpha^*) = -w(\vec{C}; \alpha)$

now : take 3-chromatic graph G with a 3-colouring α ,
and take an odd cycle C in G

- $\implies w(\vec{C}; \alpha) \neq 0$ (odd sum of +1s and -1s)

$\implies w(\vec{C}; \alpha^*) = -w(\vec{C}; \alpha) \neq w(\vec{C}; \alpha)$

$\implies \alpha$ and α^* not connected in $\mathcal{C}(G; 3)$


$\implies \mathcal{C}(G; 3)$ not connected

Mixing for larger values of $k = \chi$

- $\chi(G) = 2 \implies G$ is not 2-mixing
 - $\chi(G) = 3 \implies G$ is not 3-mixing
 - What about $k \geq 4$?
 - complete graph K_k has frozen k -colourings
- so: G has K_k as a subgraph $\implies G$ not k -mixing

Mixing for larger values of $k = \chi$

- Hajos' graph H_m ($m \geq 3$)



- has $\chi(H_m) = m$
- and is m -mixing for $m \geq 4$
- so: graphs with $k = \chi(G) \geq 4$
can be k -mixing or not k -mixing

Decision problems

k -MIXING

Input: graph G

Question: is G k -mixing?

- probably very hard, since finding one k -colouring of a graph G is probably very hard, **even if we know $k \geq \chi(G)$**

Maybe easier:

BIPARTITE- k -MIXING

Input: **bipartite** graph G

Question: is G k -mixing?

Is a given bipartite graph k -mixing ?

- trivial for $k = 2$ (“yes” if and only if G has no edges)

necessary for $k = 3$:


- for all 3-colourings α and cycles C in G : $w(\vec{C}; \alpha) = 0$

Theorem

- the condition is also **sufficient** for a graph to be 3-mixing
- so: BIPARTITE-3-MIXING is in coNP
- certificate for not 3-mixing:
3-colouring α and cycle C in G with $w(\vec{C}; \alpha) \neq 0$

A structural certificate for bipartite non-3-mixing

- pinch of two vertices at distance 2 :




- G pinchable to H : sequence of pinches changes G to H

Theorem

- connected bipartite G is not 3-mixing
 \iff G is pinchable to a chordless 6-cycle

Why the 6-cycle ?

- $C_6 \cong L_3$ – so C_6 is not 3-mixing



- **note:** C_4 is 3-mixing

Deciding bipartite mixing

- bipartite G not 3-mixing $\iff G$ pinchable to C_6

Theorem (we think)

- deciding pinchability to C_6 is NP-complete

hence

- BIPARTITE-3-MIXING is coNP-complete

Theorem

- BIPARTITE-3-MIXING is polynomial for planar graphs

open : what happens for $k \geq 4$?

A decision problem for general graphs

k -COLOUR-PATH

Input: graph G and two k -colourings α and β

Question: is there is a path in $\mathcal{C}(G; k)$ from α to β ?

or: “are α and β connected?”

- this question might be doable for any k
- trivially decidable for $k = 2$



Connected 3-colourings

- **necessary condition 1**

for two 3-colourings α and β to be connected:

- for all cycles C in G : $w(\vec{C}; \alpha) = w(\vec{C}; \beta)$


- **but not sufficient:**



Connected 3-colourings

- fixed vertex of a colouring: can never change colour


examples



- **necessary condition 2**
for two 3-colourings α and β to be connected:
 - all fixed vertices in α must be fixed in β as well
and must have the same colour in both

Connected 3-colourings

- a path P with two fixed end vertices can also be given a weight $w(\vec{P}; \alpha)$



- and this weight stays the same when recolouring
- **necessary condition 3**
for two 3-colourings α and β to be connected:
 - for all paths P with fixed ends: $w(\vec{P}; \alpha) = w(\vec{P}; \beta)$

Connected 3-colourings

- two 3-colourings α and β can only be connected if:
 - for all cycles C : $w(\vec{C}; \alpha) = w(\vec{C}; \beta)$
 - for all paths P with fixed ends : $w(\vec{P}; \alpha) = w(\vec{P}; \beta)$
 - the sets of fixed vertices in α and β must be identical

Theorem

- the conditions above are also **sufficient**
- the conditions can be checked in polynomial time

and

- if connected, then there is a path of length $O(n^2)$

k-COLOUR-PATH for $k \geq 4$

Theorem

- for $k \geq 4$, k -COLOUR-PATH is PSPACE-complete

PSPACE

- decision problems that can be solved using a **polynomial amount of memory** (no restrictions on time)
- contains **NP** and **coNP**
- equal to its **non-deterministic** variant **NPSPACE**

k -COLOUR-PATH for $k \geq 4$

Theorem

k -COLOUR-PATH for bipartite, planar graphs :

- $k = 2$: trivially decidable
- $k = 3$: decidable in polynomial time
- $k = 4$: PSPACE-complete
- $k \geq 5$: always “YES”

Length of paths between connected colourings

Theorem

- for $k \geq 4$, k -COLOUR-PATH is PSPACE-complete
- if $NP \neq PSPACE$ (similar status as $P \neq NP$), then no PSPACE-complete problem should have polynomial length certificates

so: for $k \geq 4$ path length between two connected k -colourings should not always be polynomial

Length of paths between connected colourings

Theorem

- for all $k \geq 4$, there exists graphs G with two k -colourings α and β so that
 - α and β are connected
 - the shortest path from α to β has exponential length
- the graphs can be bipartite
- and for $k = 4$ even bipartite and planar

Something different: using extra colours

- given a graph G and two k -colourings α and β
- suppose we can “buy” extra colours to go from α to β

how many extra colours do we need?

Theorem

- $\chi(G) - 1$ extra colours is always enough

$\chi - 1$ extra colours are always enough


sketch of the proof

- take a χ -colouring using colours $-1, -2, \dots, -\chi$
say with colour-classes $V_{-1}, V_{-2}, \dots, V_{-\chi}$
- starting with the k -colouring α (using colours $1, 2, \dots, k$)
 - recolour vertices in V_{-1} with colour -1
 - recolour vertices in V_{-2} with colour -2
 - *etc.*, until vertices in $V_{-(\chi-1)}$ with colour $-(\chi-1)$
- the remaining vertices in $V_{-\chi}$ form an independent set
 - hence can be recoloured to their colours according to β
- now recolour vertices in $V_{-1} \cup V_{-2} \cup \dots \cup V_{-(\chi-1)}$
according to β as well

$\chi - 1$ extra colours may be needed

Theorem

- for all C, k with $k \geq C \geq 2$
there exists graphs G with $\chi(G) = C$
and two k -colourings α and β so that
 - to get from α to β requires $C - 1$ extra colours



the graph for $C = 3$ and $k = 5$



Thank you for your attention.

