

Degrees of Perfectness

JAN VAN DEN HEUVEL

Prague, 28 July 2009

Department of Mathematics
London School of Economics and Political Science



Some good terminology

- **set system** (S, \mathcal{F}) : a finite set S
with a collection \mathcal{F} of subsets of S
- a set system is “**good**” if :
 - \mathcal{F} is closed under taking subsets, and
 - \mathcal{F} covers all of S
(for all $s \in S$ there is an $F \in \mathcal{F}$ with $x \in F$)

Two important examples

- $G = (V_G, E_G)$ be a graph
 - take \mathcal{S}_G the collection of all stable sets
(sets containing no adjacent pairs of vertices)
 - then (V_G, \mathcal{S}_G) is a good set system
- V be a vector space, and U a subset of $V \setminus \{0\}$
 - take \mathcal{I}_U the collection of all
linearly independent subsets of U
 - then (U, \mathcal{I}_U) is a good set system

Coverings

- a **covering** of (S, \mathcal{F}) :
a collection of sets from \mathcal{F} whose **union is S**
- **covering number $\text{Cov}(S, \mathcal{F})$** :
the **minimum** number of elements in a covering
- for a **graph G** : $\text{Cov}(V_G, \mathcal{S}_G)$ is the **minimum** number of
stable sets needed to **cover all vertices**
 - so $\text{Cov}(V_G, \mathcal{S}_G)$ is just the **chromatic number**

Let's make it look more complicated

- the covering number is also the solution of the IP problem :

minimise $\sum_{F \in \mathcal{F}} x_F$

subject to $\sum_{F \ni s} x_F \geq 1, \quad \text{for all } s \in S$

$$x_F \in \{0, 1, 2, \dots\}, \quad \text{for all } F \in \mathcal{F}$$

The fractional version

- removing the integrality condition :

minimise $\sum_{F \in \mathcal{F}} x_F$

subject to $\sum_{F \ni s} x_F \geq 1, \quad \text{for all } s \in S$

$$x_F \geq 0, \quad \text{for all } F \in \mathcal{F}$$

- gives the **fractional covering number** $\text{Cov}_f(S, \mathcal{F})$
 - and we obviously have : $\text{Cov}_f(S, \mathcal{F}) \leq \text{Cov}(S, \mathcal{F})$

Rule 1 of Linear Programming: dualise

- the dual LP problem of the fractional covering number is:

$$\text{maximise } \sum_{s \in S} y_s$$

$$\text{subject to } \sum_{s \in F} y_s \leq 1, \quad \text{for all } F \in \mathcal{F}$$

$$y_s \geq 0, \quad \text{for all } s \in S$$

- this gives the fractional packing number $\text{Pack}_f(S, \mathcal{F})$

- and by LP-duality: $\text{Pack}_f(S, \mathcal{F}) = \text{Cov}_f(S, \mathcal{F})$

The packing number

- the integral version is the **packing number** $\text{Pack}(S, \mathcal{F})$:
 - the maximum size $|T|$ of a subset $T \subseteq S$ so that
$$|T \cap F| \leq 1, \text{ for all } F \in \mathcal{F}$$
 - i.e.: the maximum size $|T|$ of a subset $T \subseteq S$ so that
no two elements of T appear together in a set from \mathcal{F}
- for a graph G : $\text{Pack}(V_G, \mathcal{S}_G)$ is the maximum size of
a set of vertices with no two elements in a stable set
 - so $\text{Pack}(V_G, \mathcal{S}_G)$ is just the **clique number**

The status so far

- for any good set system (S, \mathcal{F}) we have


$$\text{Pack}(S, \mathcal{F}) \leq \text{Pack}_f(S, \mathcal{F}) = \text{Cov}_f(S, \mathcal{F}) \leq \text{Cov}(S, \mathcal{F})$$

- we will add one more parameter :

the **circular covering number** $\text{Cov}_c(S, \mathcal{F})$

The circular covering number


- map the elements of S to a circle so that:
 - for every unit interval $[x, x + 1)$ along the circle elements mapped into that interval form a set from \mathcal{F}



- **circular covering number** $\text{Cov}_c(S, \mathcal{F})$:
minimum circumference of a circle for which this is possible

The right place for the circular covering number - I


- for a good set system: $\text{Cov}_c(S, \mathcal{F}) \leq \text{Cov}(S, \mathcal{F})$
 - take a disjoint cover F_1, \dots, F_k of (S, \mathcal{F})
 - put the elements of each F_i together at unit distance around a circle with circumference k :



- gives a circular cover with circumference k

The right place for the circular covering number - II

- for a good set system: $\text{Cov}_f(S, \mathcal{F}) \leq \text{Cov}_c(S, \mathcal{F})$
 - take a circular cover along a circle



- “move” the unit interval with “unit speed” round the circle
- for a set F that appears in the interval at some point:
denote by x_F the “length of time” it appears

The right place for the circular covering number - II

- for a good set system: $\text{Cov}_f(S, \mathcal{F}) \leq \text{Cov}_c(S, \mathcal{F})$
 - take a circular cover along some circle
 - for a set F that appears in the interval at some point:
denote by x_F the “length of time” it appears
- then for all $s \in S$: $\sum_{F \ni s} x_F = 1$
- and $\sum_{F \in \mathcal{F}} x_F = \text{circumference}$
- this gives a fractional cover with value the circumference

Inequalities, inequalities, and more inequalities

- so now we know :

$$\text{Pack} \leq \text{Pack}_f = \text{Cov}_f \leq \text{Cov}_c \leq \text{Cov}$$

- can we say for which good set systems we have **equality** for one of the inequalities ?
 - probably too hard (“too local”)
- what about those that satisfy an equality

“through and through” ?

$$\text{Pack} \leq \frac{\text{Cov}_f}{\text{Pack}_f} \leq \text{Cov}_c \leq \text{Cov}$$

Through and through = induced

- (S, \mathcal{F}) a good set system and $T \subseteq S$, then define:

$$\mathcal{F}_T = \{F \cap T \mid F \in \mathcal{F}\} = \{F \in \mathcal{F} \mid F \subseteq T\}$$

- then (T, \mathcal{F}_T) is again a good set system

- called an **induced** set system

- for a graph G with $U \subseteq V_G$:

$(\mathcal{S}_G)_U$ are the stable sets of the subgraph induced by U

$$\text{Pack} \leq \frac{\text{Cov}_f}{\text{Pack}_f} \leq \text{Cov}_c \leq \text{Cov}$$

Degrees of perfectness

- a good set system is **$(A = B)$ -perfect**:
 - the system and all its induced systems satisfy $A = B$
- note that we have **six** degrees of perfectness
- by definition, **perfect graphs** are exactly those graphs G
for which (V_G, \mathcal{S}_G) is **$(\text{Pack} = \text{Cov})$ -perfect**
 - that makes them **perfect** for **all inequalities** !

$$\text{Pack} \leq \frac{\text{Cov}_f}{\text{Pack}_f} \leq \text{Cov}_c \leq \text{Cov}$$

What about the other set systems ?

- we know non-perfect graphs very well :

Strong Perfect Graph Theorem (Chudnovsky et al., 2006)

- G not a perfect graph \iff

G contains an induced copy :

- of an odd cycle C_{2k+1} , $k \geq 2$, or
- of the complement $\overline{C_{2k+1}}$ of an odd cycle, $k \geq 2$

$$\text{Pack} \leq \frac{\text{Cov}_f}{\text{Pack}_f} \leq \text{Cov}_c \leq \text{Cov}$$

What about other “graphical” set systems ?

- for an odd cycle C_{2k+1} , $k \geq 2$, it is easy to check :
 - $\text{Pack}(V_{C_{2k+1}}, \mathcal{S}_{C_{2k+1}}) = 2$
 - $\text{Cov}_f(V_{C_{2k+1}}, \mathcal{S}_{C_{2k+1}}) = \text{Cov}_c(V_{C_{2k+1}}, \mathcal{S}_{C_{2k+1}}) = 2 + \frac{1}{k}$
 - $\text{Cov}(V_{C_{2k+1}}, \mathcal{S}_{C_{2k+1}}) = 3$
- similar things happen for
the complement $\overline{C_{2k+1}}$ of an odd cycle, $k \geq 2$

$$\text{Pack} \leq \frac{\text{Cov}_f}{\text{Pack}_f} \leq \text{Cov}_c \leq \text{Cov}$$

Perfect graphs are very perfect

so :

- a good set system of the form (V_G, \mathcal{S}_G) is
 - $(\text{Pack} = \text{Cov}_f)$ -perfect, or
 - $(\text{Pack} = \text{Cov}_c)$ -perfect, or
 - $(\text{Pack} = \text{Cov})$ -perfect, or
 - $(\text{Cov}_f = \text{Cov})$ -perfect, or
 - $(\text{Cov}_c = \text{Cov})$ -perfect

\iff G is perfect

$$\text{Pack} \leq \frac{\text{Cov}_f}{\text{Pack}_f} \leq \text{Cov}_c \leq \text{Cov}$$

And what about non-graphical set systems ?

- suppose (S, \mathcal{F}) is a good set system such that
 - all minimal sets not in \mathcal{F} have size 2
(smaller than 2 is not possible, as \mathcal{F} covers S)

- then form the graph G with $V_G = S$ by setting

$$s_1 s_2 \in E_G \iff \{s_1, s_2\} \notin \mathcal{F}$$

- easy to check: $(S, \mathcal{F}) = (V_G, \mathcal{S}_G)$

$$\text{Pack} \leq \frac{\text{Cov}_f}{\text{Pack}_f} \leq \text{Cov}_c \leq \text{Cov}$$

And what about non-graphical set systems ?

- (S, \mathcal{F}) is a non-graphical good set system \iff
there is a subset $T \subseteq S$ with $|T| = k \geq 3$ so that:
 - $T \notin \mathcal{F}$
 - but every proper subset of T is in \mathcal{F}
- for such a T , the induced set system (T, \mathcal{F}_T) satisfies :
 - $\text{Pack}(T, \mathcal{F}_T) = 1$
 - $\text{Cov}_f(T, \mathcal{F}_T) = \text{Cov}_c(T, \mathcal{F}_T) = 1 + \frac{1}{k-1}$
 - $\text{Cov}(T, \mathcal{F}_T) = 2$

$$\text{Pack} \leq \frac{\text{Cov}_f}{\text{Pack}_f} \leq \text{Cov}_c \leq \text{Cov}$$

Perfect graphs are really, really perfect!

so :

■ a good good set system $(\mathcal{S}, \mathcal{F})$ is

■ $(\text{Pack} = \text{Cov}_f)$ -perfect, or

■ $(\text{Pack} = \text{Cov}_c)$ -perfect, or

■ $(\text{Pack} = \text{Cov})$ -perfect, or

■ $(\text{Cov}_f = \text{Cov})$ -perfect, or

■ $(\text{Cov}_c = \text{Cov})$ -perfect

$\iff (\mathcal{S}, \mathcal{F}) = (V_G, \mathcal{S}_G)$ for some perfect graph G

$$\text{Pack} \leq \frac{\text{Cov}_f}{\text{Pack}_f} \leq \text{Cov}_c \leq \text{Cov}$$

All that is left to do . . .

- what good set systems (S, \mathcal{F}) are $(\text{Cov}_f = \text{Cov}_c)$ -perfect?
- well . . .
 - stable sets of perfect graphs
 - stable sets of odd cycles or complements of odd cycles
 - loopless matroids (vdH & Thomassé)
 - and a lot more

$$\text{Cov}_f \leq \text{Cov}_c$$

What the **** is a loopless matroid ?

- a set system (S, \mathcal{F}) is a **loopless matroid** if
 - (S, \mathcal{F}) is **good**
 - for each $F_1, F_2 \in \mathcal{F}$ with $|F_1| > |F_2|$:
there is an $s \in F_1 \setminus F_2$ so that $F_2 \cup \{s\} \in \mathcal{F}$

example

- V a **vector space**, U a subset of $V \setminus \{0\}$
- then (U, \mathcal{I}_U) is a **loopless matroid**
- so: $\text{Cov}_f(U, \mathcal{I}_U) = \text{Cov}_c(U, \mathcal{I}_U)$

$$\text{Cov}_f \leq \text{Cov}_c$$

The “remaining” case

- good set systems that are $(\text{Cov}_f = \text{Cov}_c)$ -perfect:
 - stable sets of perfect graphs
 - stable sets of odd cycles or complements of odd cycles
 - loopless matroids
 - disjoint unions of the above
 - and probably a lot more . . .

by the way :

- a stable set system (V_G, \mathcal{S}_G) is a loopless matroid

$\iff G$ is the disjoint union of cliques

$$\text{Cov}_f \leq \text{Cov}_c$$

The “remaining” case

questions :

- can we characterise $(\text{Cov}_f = \text{Cov}_c)$ -perfect set systems ?
- or at least the graphs G for which (V_G, \mathcal{S}_G) is
 $(\text{Cov}_f = \text{Cov}_c)$ -perfect ?
- what “natural” class of set systems
contains both matroids and stable sets of perfect graphs ?

$$\text{Cov}_f \leq \text{Cov}_c$$