

Turán's theorem for pseudo-random graphs

Yoshiharu Kohayakawa^{a,1}, Vojtěch Rödl^{b,2}, Mathias Schacht^{c,3},
Papa Sissokho^d, Jozef Skokan^{a,4}

^a Instituto de Matemática e Estatística, Universidade de São Paulo, Rua do Matão 1010, 05508-090 São Paulo, Brazil

^b Department of Mathematics and Computer Science, Emory University, Atlanta, GA 30322, USA

^c Humboldt-Universität zu Berlin, Institut für Informatik, Unter den Linden 6, D-10099 Berlin, Germany

^d Mathematics Department, Illinois State University, Campus Box 4520, Stevenson Hall 313,
Normal, IL 61790-4520, USA

Received 18 December 2005

Available online 16 January 2007

Dedicated to Professor R.L. Graham on the occasion of his 70th birthday

Abstract

The generalized Turán number $\text{ex}(G, H)$ of two graphs G and H is the maximum number of edges in a subgraph of G not containing H . When G is the complete graph K_m on m vertices, the value of $\text{ex}(K_m, H)$ is $(1 - 1/(\chi(H) - 1) + o(1))\binom{m}{2}$, where $o(1) \rightarrow 0$ as $m \rightarrow \infty$, by the Erdős–Stone–Simonovits theorem.

In this paper we give an analogous result for triangle-free graphs H and pseudo-random graphs G . Our concept of pseudo-randomness is inspired by the *jumbled* graphs introduced by Thomason [A. Thomason, Pseudorandom graphs, in: Random Graphs '85, Poznań, 1985, North-Holland, Amsterdam, 1987, pp. 307–331. MR 89d:05158]. A graph G is (q, β) -*bi-jumbled* if

$$|e_G(X, Y) - q|X||Y|| \leq \beta \sqrt{|X||Y|}$$

for every two sets of vertices $X, Y \subset V(G)$. Here $e_G(X, Y)$ is the number of pairs (x, y) such that $x \in X$, $y \in Y$, and $xy \in E(G)$. This condition guarantees that G and the binomial random graph with edge probability q share a number of properties.

E-mail addresses: yoshi@ime.usp.br (Y. Kohayakawa), rodl@mathcs.emory.edu (V. Rödl), schacht@informatik.hu-berlin.de (M. Schacht), psissok@ilstu.edu (P. Sissokho), skokan@ime.usp.br (J. Skokan).

¹ The author was partially supported by FAPESP and CNPq through a Temático-ProNEx project (Proc. FAPESP 2003/09925-5) and by CNPq (Proc. 306334/2004-6 and 479882/2004-5).

² The author was partially supported by NSF Grant DMS 0300529.

³ The author was supported by DFG Grant SCHA 1263/1-1.

⁴ The author was supported by NSF Grant INT-0305793, by NSA Grant H98230-04-1-0035, by CNPq (Proc. 479882/2004-5), and by FAPESP (Proj. Temático-ProNEx Proc. FAPESP 2003/09925-5 and Proc. FAPESP 2004/15397-4).

Our results imply that, for example, for any triangle-free graph H with maximum degree Δ and for any $\delta > 0$ there exists $\gamma > 0$ so that the following holds: any large enough m -vertex, $(q, \gamma q^{\Delta+1/2}m)$ -bi-jumbled graph G satisfies

$$\text{ex}(G, H) \leq \left(1 - \frac{1}{\chi(H) - 1} + \delta\right) |E(G)|.$$

© 2006 Elsevier Inc. All rights reserved.

Keywords: Turán's theorem; Pseudo-randomness; Regularity lemma; (n, d, λ) -Graphs

1. Introduction

We say that a graph is H -free if it does not contain a copy of a given graph H as a subgraph (not necessarily induced). A classical area of extremal graph theory investigates numerical and structural results concerning H -free graphs. A basic problem in this area is to determine, or estimate, the maximum number of edges $\text{ex}(m, H)$ that an H -free graph on m vertices may have. When H is a complete graph, we know the value of $\text{ex}(m, H)$ precisely, by Turán's theorem [34] (hence we refer to $\text{ex}(m, H)$ as the *Turán number of H*). When H is arbitrary, an asymptotic solution to this problem is given by the celebrated Erdős–Stone–Simonovits theorem, at least when $\chi(H) \geq 3$.

Theorem 1. (Erdős, Stone, Simonovits [10,12]) *For every graph H with chromatic number $\chi(H)$,*

$$\text{ex}(m, H) = \left(1 - \frac{1}{\chi(H) - 1} + o(1)\right) \binom{m}{2}, \quad (1)$$

where $o(1) \rightarrow 0$ as $m \rightarrow \infty$.

Here we are interested in a variant of the function $\text{ex}(m, H)$. Denote by $\text{ex}(G, H)$ the maximum number of edges that an H -free subgraph of a given graph G may have, i.e.,

$$\text{ex}(G, H) = \max\{|E(G')| : H \not\subset G' \subset G\}.$$

For instance, if $G = K_m$, the complete graph on m vertices, then $\text{ex}(K_m, H)$ is the usual Turán number $\text{ex}(m, H)$. Furthermore, by considering a random partition of vertices of G into $\chi(H) - 1$ parts, one easily observes that

$$\text{ex}(G, H) \geq \left(1 - \frac{1}{\chi(H) - 1}\right) |E(G)| \quad (2)$$

holds for any G and H .

Let us mention a few problems and results concerning the “generalized Turán function” $\text{ex}(G, H)$. The case in which G is the n -dimensional hypercube Q^n and H is a short even cycle was raised by Erdős in [8], and the best results in this direction are due to Chung [4].

Two results for the case in which G is the random graph $\mathcal{G}(m, q)$ and $H = K_3$ are due to Frankl and Rödl [13] and Babai, Simonovits, and Spencer [3]. In [14], Füredi investigates $\text{ex}(\mathcal{G}(m, q), C_4)$ (see also [19]). For detailed discussions on the function $\text{ex}(\mathcal{G}(m, q), H)$, the reader is referred to [15], [17, Chapter 8], [21], and [23], and the references therein.

Roughly speaking, the main problem is to identify the threshold for $q = q(m)$ for the property that the lower bound in (2) should be asymptotically tight, giving the almost sure value of $\text{ex}(\mathcal{G}(m, q), H)$. The original conjecture about the threshold for H arbitrary, which is still open, may be found in [20].

For deterministic graphs G the known results in this direction are for the so-called (m, d, λ) -graphs. Let G be a graph and let $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m$ be the eigenvalues of its adjacency matrix. We say that G is an (m, d, λ) -graph if it has m vertices, it is d -regular and $\max\{\lambda_2, -\lambda_m\} \leq \lambda$. Sudakov, Szabó, and Vu [31] proved that, for $t \geq 3$,

$$\text{ex}(G, K_t) = \left(1 - \frac{1}{t-1} + o(1)\right) |E(G)| \tag{3}$$

for any (m, d, λ) -graph G , as long as

$$d^{t-1}/m^{t-2} \gg \lambda, \tag{4}$$

that is, $\lim_{m \rightarrow 0} m^{t-2}\lambda/d^{t-1} = 0$. A result of Krivelevich, Sudakov, and Szabó [29, Theorem 1.2], inspired by a construction of Alon [1], implies that condition (4) is essentially best possible for $t = 3$ and basically any $d = d(n) = \Omega(n^{2/3})$. For $t > 3$ it is not known whether condition (4) is optimal.

Recently, Chung [5] obtained a result analogous to the one of Sudakov, Szabó, and Vu above, applicable to non-regular graphs G , by considering the spectrum of normalized Laplacians.

In this paper, we prove a result similar to the ones of Sudakov, Szabó, and Vu and of Chung, with K_t replaced by arbitrary triangle-free graphs H and certain ‘pseudo-random’ graphs G . First, we need to define some simple graph parameters. For any graph H , we define the *degeneracy* d_H of H by

$$d_H = \max\{\delta(H'): H' \subset H\},$$

where $\delta(H')$ denotes the minimum degree of the graph H' . We remark that d_H is the smallest integer d for which there exists an ordering u_1, \dots, u_h of the vertices of H ($h = |V(H)|$) in which u_i has at most d neighbors among u_1, \dots, u_{i-1} , for every $1 \leq i \leq h$. Any such ordering is called a *d-degenerate ordering*.

Moreover, let

$$D_H = \min\{2d_H, \Delta(H)\},$$

where $\Delta(H)$ stands for the maximum degree of H . We shall later make use of the following simple fact: for any d_H -element set F of vertices of H , there is a D_H -degenerate ordering u_1, \dots, u_h of $V(H)$ with $F = \{u_1, \dots, u_{d_H}\}$. Finally, let

$$v_H = \frac{1}{2}(d_H + D_H + 1). \tag{5}$$

Remark 2. The parameters D_H and v_H are somewhat artificial. For simplicity, the reader may prefer to replace D_H by $\Delta(H)$ in (5) at the first reading, although some statements below are considerably weaker with this change. For example, if $H = K_{2,t}$ and t is large, then $d_H = 2$, $D_H = 4$, $v_H = 7/2$, and condition (6) becomes much more restrictive with v_H replaced by $(d_H + \Delta(H) + 1)/2 = (t + 3)/2 \gg 7/2$.

Our main theorem, to be given in a short while, implies the following result for (m, d, λ) -graphs.

Theorem 3. *Let a triangle-free graph H and $\delta > 0$ be given. Then there exists $\gamma = \gamma(\delta, H) > 0$ such that for any function $d = d(m) < m$ there is $M_0 = M_0(\delta, H, d)$ such that any (m, d, λ) -graph G with $m \geq M_0$ vertices and satisfying*

$$\gamma d^{v_H} / m^{v_H-1} \geq \lambda \tag{6}$$

has

$$\text{ex}(G, H) \leq \left(1 - \frac{1}{\chi(H) - 1} + \delta\right) |E(G)|.$$

Notice that to satisfy (6) for graphs with $\lambda = O(\sqrt{d})$ (e.g., for Ramanujan graphs), it suffices to have $d^{2v_H-1} = d^{D_H+d_H} \gg m^{D_H+d_H-1} = m^{2v_H-2}$. Theorem 3 is a consequence of a more general theorem for pseudo-random graphs; see Theorem 5 in the next section.

This paper is organized as follows. In the next section, we state and discuss Theorem 5, as well as derive Theorem 3 from it. In Section 3, we present additional definitions and notation and give a fairly detailed outline of the proof of Theorem 5. In Section 3 we also state all the auxiliary lemmas (including what we call the Embedding Lemma, the Regularity-to-Pair Lemma, and the Pair-to-Tuple Lemma) needed for the proof of Theorem 5. In Section 4 we give the proof of Theorem 5 and in Section 5 we prove some technical facts needed in the proofs of the Embedding Lemma and the Pair-to-Tuple Lemma. Sections 6 and 7 contain the proofs of the Embedding Lemma and the Pair-to-Tuple Lemma.

2. The result for pseudo-random graphs

A graph is *pseudo-random* if it resembles (in some well-defined sense) a random graph of the same density. The systematic study of such graphs was initiated by Thomason [33], who introduced the notion of *jumbled* graphs. A graph G is (q, β) -*jumbled* if for every $X \subset V(G)$ we have

$$\left| e_G(X) - q \binom{|X|}{2} \right| \leq \beta |X|,$$

where $e_G(X)$ denotes the number of edges in G with both endpoints in X .

Here we shall use a closely related concept of pseudo-randomness. For any sets $X, Y \subset V$, we write

$$E_G(X, Y) = \{(x, y): x \in X, y \in Y, \{x, y\} \in E(G)\}.$$

We also set $e_G(X, Y) = |E_G(X, Y)|$ and $d_G(X, Y) = e_G(X, Y)/|X||Y|$. Note that each edge in $X \cap Y$ is counted twice in $e_G(X, Y)$. Thus, if $X = Y$ then $e_G(X) = e_G(X, X)/2$ is the number of edges in G with both endpoints in X . We drop the subscript G whenever there is no danger of confusion.

Definition 4. We say that a graph G is (q, β) -*bi-jumbled* if for every $X, Y \subset V(G)$ we have

$$|e_G(X, Y) - q|X||Y|| \leq \beta \sqrt{|X||Y|}. \tag{7}$$

It is easy to see that every (q, β) -bi-jumbled graph is also $(q, (\beta + q)/2)$ -jumbled. Since we consider only $0 \leq q \leq 1 < \beta$, we immediately have that every (q, β) -bi-jumbled graph is (q, β) -jumbled.

To put (7) into some context, on the one hand, we observe that the random graph $\mathcal{G}(m, q)$ is almost surely (q, β) -bi-jumbled for $\beta = O(\sqrt{qm})$ if, say, $qm \gg \log m$.⁵ On the other hand, Erdős and Spencer [11] (see also Theorem 5 in [9]) observed that there exists $c > 0$ such that every m -vertex graph with density q contains two disjoint sets X and Y for which $|e(X, Y) - q|X||Y|| \geq c\sqrt{qm}\sqrt{|X||Y|}$, as long as $q(1 - q) \geq 1/m$.

We may finally state our main theorem, Theorem 5. Our main result shows that if β is sufficiently small, then (2) is asymptotically optimal for any sufficiently large (q, β) -bi-jumbled graph G and any triangle-free graph H .

Theorem 5. *Let a triangle-free graph H and $\delta > 0$ be given. Then there exists $\gamma = \gamma(\delta, H) > 0$ with the following property: For any function $q = q(m)$ there is $M_0 = M_0(\delta, H, q)$ such that any $(q, \gamma q^{v_H} m)$ -bi-jumbled graph G with $m \geq M_0$ vertices has*

$$\text{ex}(G, H) \leq \left(1 - \frac{1}{\chi(H) - 1} + \delta\right) |E(G)|. \tag{8}$$

Remark 6. From $c\sqrt{qm} \leq \beta = \gamma q^{v_H} m$ we deduce that the inequality $q \geq q_{\gamma, H}(m) := (c^{-2}\gamma^2 m)^{-1/(2v_H - 1)}$ must hold for any m -vertex, $(q, \gamma q^{v_H} m)$ -bi-jumbled graph. Hence, we may assume that the function q in Theorem 5 satisfies $q \geq q_{\gamma, H}(m)$ for otherwise there is no $(q, \gamma q^{v_H} m)$ -bi-jumbled graph G and the statement of Theorem 5 holds trivially.

Theorem 5 gives a sufficient condition on the bi-jumbledness of G for (8) to hold. Assuming the best possible bi-jumbledness $\beta = O(\sqrt{qm})$, the bi-jumbledness hypothesis $\beta = \gamma q^{v_H} m$ in Theorem 5 gives a condition on the density q for (8) to hold. We illustrate this on the concrete example $H = C_5$. We have $v_{C_5} = 5/2$. Theorem 5 implies that, for any $(q, O(\sqrt{qm}))$ -bi-jumbled graph G , we have $\text{ex}(G, C_5) = (1/2 + o(1))|E(G)|$ if $q^{5/2} m \gg \sqrt{qm}$, that is, $q \gg m^{-1/4}$. However, a fairly simple argument based on the Sparse Regularity Lemma (Proposition 9) yields that the much weaker condition $q \gg m^{-1/2}$ actually suffices.

Let us mention that there exists a C_5 -free, $(q, O(\sqrt{qm}))$ -bi-jumbled graph A_5 with $q = \Theta(m^{-3/5})$. Thus, the “threshold” for this problem lies between $m^{-3/5}$ and $m^{-1/2}$. Hence, even in this case, we do not have an optimal result. For a general odd cycle $C_{2\ell+1}$, the “threshold” is between $m^{-1+2/(2\ell+1)}$ and $m^{-1+1/\ell}$. The lower end of the gap is proved considering a $C_{2\ell+1}$ -free, $(q, O(\sqrt{qm}))$ -bi-jumbled graph $A_{2\ell+1}$ with $q = \Theta(m^{-1+2/(2\ell+1)})$. The existence of the graphs $A_{2\ell+1}$ may be proved suitably adapting a beautiful construction of Alon [1] (see also [28, Section 3, Example 10]).

We believe that it would be interesting to weaken the bi-jumbledness hypothesis in Theorem 5. Moreover, it would be interesting to drop the triangle-freeness condition on the graph H . In this direction, we only mention that in [25] a result similar to Theorem 5 for arbitrary graphs H is proved, but a stronger bi-jumbledness hypothesis on G is required. We finish this section deducing Theorem 3 from Theorem 5.

⁵ This fact may be checked by combining Lemma 3.8 in [16] and the fact that, almost surely, all vertices of $\mathcal{G}(m, q)$ have degree $(1 + o(1))qm$ for $qm \gg \log m$.

Proof of Theorem 3. It is a well-known fact (see Corollary 9.2.5 in [2]) that any (m, d, λ) -graph G is $(d/m, \lambda)$ -bi-jumbled. Therefore, the graph G will be $(d/m, \gamma(d/m)^{\nu_H} m)$ -bi-jumbled if

$$\lambda \leq \gamma(d/m)^{\nu_H} m, \tag{9}$$

which is equivalent to (6). \square

3. An outline of the proof of Theorem 5 and auxiliary results

In this section we introduce all the necessary tools for the proof of Theorem 5. We also try to motivate these tools by discussing the underlying ideas in the proof.

3.1. Additional definitions and notation

We start with some basic notation. Let ℓ be a positive integer. We denote by $[\ell]$ the set $\{1, 2, \dots, \ell\}$. For a multiset $I = \{i_1, \dots, i_r\}$ we write $I \subset [\ell]$ to mean that $i_1, \dots, i_r \in [\ell]$. We also adopt the convention that we always write the elements of I in non-decreasing order, i.e., $i_1 \leq \dots \leq i_r$. For three real numbers a, b , and c , the expression $a = b \pm c$ means $b - c \leq a \leq b + c$. We also write a/bc instead of $a/(bc)$ whenever there is no danger of confusion. For functions $f = f(n)$ and $g = g(n)$ we write $f \gg g$ and $g \ll f$ if $\lim_{n \rightarrow \infty} f(n)/g(n) = +\infty$. For clarity, we omit inessential floor and ceiling brackets.

Let $G = (V, E)$ be a graph. For a vertex $x \in V$ let $N(x)$ be the set of all neighbors of x in G . If $U \subset V$ then $N_U(x)$ denotes the set of neighbors of $x \in V$ belonging to U , that is, $N_U(x) = N(x) \cap U$. For an r -set $X = \{x_1, \dots, x_r\} \subset V$ and a set $U \subset V$, we let $N(X) = N(x_1, \dots, x_r) = N(x_1) \cap \dots \cap N(x_r)$ and $N_U(X) = N_U(x_1, \dots, x_r) = N(x_1) \cap \dots \cap N(x_r) \cap U$.

We say that J is an (ℓ, n, p) -partite graph if J is ℓ -partite with $V(J) = \bigcup_{j=1}^{\ell} V_j$, $|V_j| = n$ for all $j \in [\ell]$, and $e(J[V_i, V_j]) = pn^2$ for all $i \neq j \in [\ell]$.

For an (ℓ, n, p) -partite graph J , an integer $r \geq 1$, and a multiset $I = \{i_1, \dots, i_r\} \subset [\ell]$, denote by $\mathcal{T}(I)$ the set of all r -tuples $(x_1, \dots, x_r) \in V_{i_1} \times \dots \times V_{i_r}$ such that $x_i \neq x_j$ for all $1 \leq i < j \leq r$. Note that

$$(n - r)^r < (n - r + 1)^r \leq |\mathcal{T}(I)| \leq n^r, \tag{10}$$

since each x_i can be chosen in at least $n - r + 1$ ways to avoid x_1, \dots, x_{i-1} .

Now we define two important properties of (ℓ, n, p) -partite graphs.

Definition 7. An (ℓ, n, p) -partite graph J has property $\text{TUPLE}_{\ell}(\varepsilon, d)$ if for every integer $1 \leq r \leq d$, every multiset $I = \{i_1, \dots, i_r\} \subset [\ell]$, and for all $j \in [\ell] \setminus I$, we have

$$|N_{V_j}(x_1, \dots, x_r)| = (1 \pm \varepsilon)p^r n$$

for all but at most εn^r r -tuples $(x_1, \dots, x_r) \in \mathcal{T}(I)$.

When $d = 2$ we say that J satisfies the pair condition $\text{PAIR}_{\ell}(\varepsilon)$.

3.2. An outline of the proof of Theorem 5

We now outline the proof of Theorem 5. We hope that this will motivate the somewhat technical looking auxiliary lemmas that will be required. Our proof strategy is natural and proceeds as follows: consider an arbitrary spanning subgraph G' of G with

$$|E(G')| \geq \left(1 - \frac{1}{\chi(H) - 1} + \delta\right) |E(G)|. \tag{11}$$

Suppose first that the density q of G is constant, independent of n . We apply Szemerédi’s regularity lemma [32] with ε significantly smaller than δ and q , and obtain a partition $\bigcup_{j=0}^t V_j$ of the vertex set of G' into a bounded number $t + 1$ of parts so that all but at most $\varepsilon \binom{t}{2}$ pairs (V_i, V_j) , $1 \leq i < j \leq t$, are ε -regular, i.e., for all $V'_i \subset V_i$ and $V'_j \subset V_j$ with $|V'_i| \geq \varepsilon |V_i|$ and $|V'_j| \geq \varepsilon |V_j|$, we have

$$|d_G(V_i, V_j) - d_G(V'_i, V'_j)| \leq \varepsilon.$$

Standard arguments (see, e.g., [7, Section 7.5, pp. 186–187]) show that there exist $\ell = \chi(H)$ sets (without loss of generality, V_1, \dots, V_ℓ) so that there is an (ℓ, n, p) -partite graph $J \subset G'$ with ℓ -partition $\bigcup_{j=1}^\ell V_j$ such that $p = \alpha q$, α is considerably larger than ε , and each pair (V_i, V_j) is ε -regular.⁶ It is a well-known fact (e.g., Theorem 3.1 in [27]) that J satisfying the conditions above also contains a copy of H . We remark that this statement (sometimes called the Embedding Lemma) is ensured by the regularity of J itself.

For the case $q = o(1)$ we utilize the same approach. First we apply a version of the regularity lemma for sparse graphs (Proposition 9) to G' and obtain an (ε, G', q) -regular partition $V_0 \cup V_1 \cup \dots \cup V_t$ of $V(G')$ (see Section 3.4 for the relevant definitions). In the same way as above (see Section 4 for the details) we obtain an (ℓ, n, p) -partite graph J with ℓ -partition $\bigcup_{j=1}^\ell V_j$ with $p = \alpha q$ for some constant $\alpha > 0$ and such that each pair (V_i, V_j) is (ε, q) -regular. Unfortunately, the embedding lemma required in this context does not hold (see, e.g., [21, Theorem B’]).

One way to deal with this problem is to restrict the choice of G to certain classes of graphs (such as random graphs) and to prove an appropriate embedding lemma that works for their subgraphs G' (for instance, see Theorem B’’ in [21] and Lemmas 2.2 and 2.2’ in [23]). In this paper, roughly speaking, we follow an approach in [26]. An embedding lemma in [26] is as follows:

(*) *Let H be a triangle-free graph and C a positive constant. If J is an n -vertex graph with density $p = p(n) = |E(J)| \binom{n}{2}^{-1} \gg n^{-1/D_H}$ satisfying properties BDD(C, D_H) and PAIR given below, then J contains H as a subgraph, as long as n is sufficiently large.*

BDD(C, D_H): $|N_J(x_1, \dots, x_r)| \leq C^r p^r n$ holds for all mutually distinct vertices $x_1, \dots, x_r \in V(J)$ and $1 \leq r \leq D_H$.

PAIR: $|N_J(x_1, x_2)| = (1 + o(1)) p^2 n$ holds for all but at most $o(n^2)$ pairs $\{x_1, x_2\} \subset V(J)$.

⁶ From $\bigcup_{j=0}^t V_j$ we construct a “cluster” graph F_c on $[t]$ whose edges are those pairs $\{i, j\}$ for which the pair (V_i, V_j) is ε -regular and has large density. The graph F_c has enough edges for us to apply Turán’s theorem and obtain a copy of $K_{\chi(H)}$. To deduce this we need $d_{G'}(V_i, V_j) \leq (1 + o(1))q$ for all $i \neq j$, which may be guaranteed by the bi-jumbledness of G . This copy of $K_{\chi(H)}$ corresponds to a subgraph of G' that may be further reduced or “sliced” (see Lemma 10) to the $(\chi(H), n, p)$ -partite graph J we are looking for.

Going back to the proof of Theorem 5, we recall that we had arrived at an (ℓ, n, p) -partite graph $J \subset G'$ with ℓ -partition $\bigcup_{j=1}^{\ell} V_j$ such that $p = \alpha q$ with α a positive constant and each pair (V_i, V_j) is (ε, q) -regular. The first discrepancy that one notices between our current setup and the hypotheses in (*) is that our J is ℓ -partite, whereas in (*) we do not have an ℓ -partite graph. As the reader may guess, albeit cumbersome, this difference is not essential, and one may in fact prove an appropriate “ ℓ -partite version” of (*). In the discussion that follows, for simplicity, when not important, we shall blur this discrepancy and we shall ignore the fact that we have an ℓ -partite graph J at hand.

A more substantial discrepancy occurs in the hypotheses BDD(C, D_H) and PAIR in (*): in our proof of Theorem 5, we have arrived at an ℓ -partite (ε, q) -regular graph J .

Achieving PAIR

The reader may be familiar with the fact that in the case of dense graphs, the $o(1)$ -regularity of a pair (V_i, V_j) and property PAIR($o(1)$) are equivalent (in a certain precise sense, see [6,33] for details). Unfortunately, this equivalence breaks down in the sparse setting, as observed in Theorem A' in [21]. Therefore, achieving hypothesis PAIR in (*) requires some work. This will be accomplished by making use of Proposition 11 below, which, roughly speaking, states that one recovers the fact that $o(1)$ -regularity implies PAIR if one has a graph J that is a subgraph of a suitably bi-jumbled graph Γ , as long as one has a positive fraction α of the edges of Γ in J .

Loosening BDD

Let us now discuss hypothesis BDD(C, D_H) in (*). Basically, if G has property BDD(C, D_H), then any subgraph $J \subset G$ with a positive fraction of the edges of G has BDD(C', D_H) for some $C' \geq C$. As it turns out, the constant C' ends up depending on some other parameters in the proof in such a way that we are not able to use this simple hereditary property of BDD. Therefore, we take a different route. For every $1 \leq r \leq D_H$, define an r -uniform hypergraph \mathcal{B}_r on the vertex set of J , putting an r -set $B \subset V(J)$ in \mathcal{B}_r if the joint neighborhood $N_J(B)$ of B in J violates the upper bound in the definition of BDD(C, D_H), that is, $|N_J(B)| > C' p^r n$. A simple consequence of the bi-jumbledness of G is that the hypergraphs \mathcal{B}_r are in a certain sense locally sparse: for every $1 \leq r \leq D_H$, if an $(r-1)$ -set is not a member of \mathcal{B}_{r-1} , then it cannot be contained in many members of \mathcal{B}_r (see Lemma 14). This sparseness of the \mathcal{B}_r turns out to be enough for our purposes.

The Embedding Lemma

As the discussion above suggests, the embedding lemma that we shall make use of is an ℓ -partite variant of (*), with the BDD hypothesis replaced by the hypothesis that J should be a subgraph of a suitably bi-jumbled graph G , with a positive fraction α of the edges of G in J . From this hypothesis, one obtains PAIR and the local sparseness of the \mathcal{B}_r . For the precise statement of this embedding lemma, see Proposition 8.

Before we finish this outline, we just mention a step in the proof of this embedding lemma. We remark that we shall use the bi-jumbledness of G to show that, in fact, PAIR implies the following property:

TUPLE(d_H): $|N_J(x_1, \dots, x_r)| = (1 + o(1))p^r n$ holds for all but at most $o(n^r)$ r -sets $\{x_1, \dots, x_r\} \subset V(J)$ for any $1 \leq r \leq d_H$

(see Proposition 12). Going from PAIR to TUPLE(d_H) is also an important step in the proof of (*). For this step, hypothesis BDD is used in [26]; here, in the proof of Proposition 12, we again replace BDD with the local sparseness of the \mathcal{B}_r .

3.3. An embedding lemma for ℓ -partite bi-jumbled graphs

In this section we state the adjusted version of one of the main results from [26] (see (*)) discussed in Section 3.2. Given an ℓ -partite graph H with ℓ -partition $V(H) = \bigcup_{j=1}^{\ell} U_j$, an embedding of H in an (ℓ, n, p) -partite graph J is an injective, edge preserving map $f: V(H) \rightarrow V(J)$ such that $f(U_j) \subset V_j$ for all $1 \leq j \leq \ell$. The next proposition shows that TUPLE $_{\ell}$ and large enough density guarantee an embedding of any triangle-free graph H in J , as long as J satisfies a certain sparseness condition (see (12) below).

Proposition 8 (Embedding Lemma). *Let H be a fixed triangle-free, ℓ -partite graph with h vertices and e edges. Then for all $0 < \alpha, \eta \leq 1$ there exist $\varepsilon = \varepsilon(H, \alpha, \eta) > 0$ and $\gamma = \gamma(H, \alpha, \eta) > 0$ such that for any function $p = p(n)$ satisfying $p^{d_H} n \gg 1$ there is $N_1 = N_1(H, \alpha, \eta, p) > 0$ for which the following holds. Suppose that*

- (a) J is an (ℓ, n, p) -partite graph and $n > N_1$,
- (b) for all $U \subset V_i$ and $W \subset V_j$, $i \neq j \in [\ell]$, we have

$$e_J(U, W) \leq \frac{p}{\alpha} |U||W| + \gamma \left(\frac{p}{\alpha}\right)^{v_H} n \sqrt{|U||W|}, \tag{12}$$

- (c) J satisfies TUPLE $_{\ell}(\varepsilon, d_H)$.

Then the number of embeddings of H in J is at least $(1 - \eta)p^e n^h$.

We shall apply Proposition 8 to an (ℓ, n, p) -partite graph J that is obtained from a subgraph G' of a $(q, \gamma q^{v_H} m)$ -bi-jumbled graph G , as explained in Section 3.2. Moreover, we shall have $p = \alpha q$ for some constant $\alpha > 0$. Condition (12) will follow from the upper bound in the $(q, \gamma q^{v_H} m)$ -bi-jumbledness hypothesis on G (see (7)), by substituting p/α for q . The proof of Proposition 8 is given in Section 6.

3.4. The Sparse Regularity Lemma

Let $G = (V, E)$ be a graph. Suppose $0 < q \leq 1$, $\xi > 0$ and $C > 1$. For two disjoint subsets X, Y of V , we let

$$d_{G,q}(X, Y) = \frac{e_G(X, Y)}{q|X||Y|},$$

which we refer to as the q -density of the pair (X, Y) .

We say that G is a (ξ, C) -bounded graph with respect to density q if for all pairwise disjoint $X, Y \subset V$, with $|X|, |Y| \geq \xi|V|$, we have $e_G(X, Y) \leq Cq|X||Y|$.

For $\varepsilon > 0$ fixed and $X, Y \subset V$, $X \cap Y = \emptyset$, we say that the pair (X, Y) is (ε, q) -regular if for all $X' \subset X$ and $Y' \subset Y$ with

$$|X'| \geq \varepsilon|X| \quad \text{and} \quad |Y'| \geq \varepsilon|Y|,$$

we have

$$|d_{G,q}(X, Y) - d_{G,q}(X', Y')| \leq \varepsilon.$$

Note that for $q = 1$ we get the well-known definition of ε -regularity [32].

When G is (ℓ, n, p) -partite with ℓ -partition $\bigcup_{i=1}^{\ell} V_i$ we say that G is (ε, q) -regular if all pairs (V_i, V_j) , $1 \leq i < j \leq \ell$, are (ε, q) -regular.

Let $\bigcup_{j=0}^t V_j$ be a partition of V . We call V_0 the *exceptional class*. This partition is called (ε, t) -equitable if $|V_0| \leq \varepsilon|V|$ and $|V_1| = \dots = |V_t|$.

We say that an (ε, t) -equitable partition $\bigcup_{j=0}^t V_j$ of V is (ε, G, q) -regular if all but at most $\varepsilon \binom{t}{2}$ pairs (V_i, V_j) , $1 \leq i < j \leq k$, are (ε, q) -regular. Now we can state a variant of Szemerédi’s regularity lemma [32] for sparse graphs (see, e.g., [18,22]).

Proposition 9 (*Sparse Regularity Lemma*). *For any $\varepsilon > 0$, $C > 1$, and $t_1 \geq 1$, there exist constants $T_1 = T_1(\varepsilon, C, t_1)$, $\xi = \xi(\varepsilon, C, t_1) \leq \min\{1/2T_1, \varepsilon\}$, and $M_1 = M_1(\varepsilon, C, t_1)$ such that any graph G with at least M_1 vertices that is (ξ, C) -bounded with respect to density $0 < q \leq 1$ admits an (ε, t) -equitable (ε, G, q) -regular partition of its vertex set with $t_1 \leq t \leq T_1$.*

After applying the lemma above we obtain (ε, q) -regular bipartite graphs with different densities. The next lemma will allow us to change these densities to a particular value without losing regularity.

Lemma 10 (*Slicing Lemma* [23]). *For every $0 < \alpha$, $\varepsilon \leq 1$, $C > 1$, and a function $q = q(n)$ satisfying $qn \gg 1$ there exists $n_0 = n_0(\alpha, \varepsilon, C, q)$ such that if $B = (U \cup W, E)$ is a bipartite graph satisfying*

- (i) $|U| = |W| = n > n_0$,
- (ii) $\alpha q|U||W| \leq e_B(U, W) \leq Cq|U||W|$, and
- (iii) B is (ε, q) -regular,

then there exists a $(3\varepsilon, q)$ -regular subgraph $B' = (U \cup W, E') \subset B$ such that $e_{B'}(U, W) = \alpha q|U||W|$.

3.5. Regularity and the pair condition

The next proposition shows that, under certain restrictions, a regular (ℓ, n, p) -partite graph also has property PAIR_ℓ .

Proposition 11 (*Regularity-to-Pair Lemma* [25]). *For any $0 < \alpha$, $\varrho \leq 1$ and any integer $\ell > 1$, there exist $\delta = \delta(\alpha, \varrho, \ell) > 0$ and $\gamma = \gamma(\alpha, \varrho, \ell) > 0$ such that for every function $q = q(n)$ there exists an $n_0 = n_0(\alpha, \varrho, \ell, q) > 1$ for which the following holds.*

Let Γ be a $(q, \gamma q^2 \ell n)$ -bi-jumbled graph on ℓn vertices with $n > n_0$. Suppose J is a (δ, q) -regular (ℓ, n, p) -partite subgraph of Γ satisfying $p \geq \alpha q$. Then J also has property $\text{PAIR}_\ell(\varrho)$.

We remark that if p is a constant then the above statement holds for any $(\delta, 1)$ -regular (ℓ, n, p) -partite graph J (i.e., J need not to be a subgraph of some bi-jumbled graph Γ).

Clearly, any graph having property $\text{TUPLE}_\ell(\varepsilon, d)$, $d \geq 2$, also satisfies $\text{PAIR}_\ell(\varepsilon)$. The next proposition shows that under certain conditions the converse is also true.

Proposition 12 (Pair-to-Tuple Lemma). *Let $d \geq 1$ and $\ell \geq 2$ be integers. Then for every $0 < \alpha, \varepsilon \leq 1$ there exist $\delta = \delta(d, \ell, \alpha, \varepsilon) > 0$ and $\gamma = \gamma(d, \ell, \alpha, \varepsilon) > 0$ such that for every function $p = p(n)$ satisfying $p^d n \gg 1$ there is $N_0 = N_0(d, \ell, \alpha, \varepsilon, p)$ with the following property: any (ℓ, n, p) -partite graph J with $n > N_0$ satisfying*

(i) *for all $U \subset V_i$ and $W \subset V_j, i \neq j \in [\ell]$, we have*

$$e_J(U, W) \leq \frac{p}{\alpha} |U||W| + \gamma \left(\frac{p}{\alpha}\right)^{(d+3)/2} n \sqrt{|U||W|}, \tag{13}$$

(ii) $\text{PAIR}_\ell(\delta)$,

also satisfies $\text{TUPLE}_\ell(\varepsilon, d)$.

We shall apply Proposition 12 with $d = d_H$ to an (ℓ, n, p) -partite graph J that is obtained from a subgraph G' of a $(q, \gamma q^{(d_H+3)/2} m)$ -bi-jumbled graph G , as explained in Section 3.2. Moreover, we shall have $p = \alpha q$ for some constant $\alpha > 0$. Condition (13) will follow from the upper bound in the $(q, \gamma q^{(d_H+3)/2} m)$ -bi-jumbledness hypothesis on G (see (7)), by substituting p/α for q . The proof of Proposition 12 appears in Section 7.

4. Proof of the main result

Now we are ready to prove the main result, Theorem 5. When $D_H = 1$, the graph H is a matching and we just need $qm > (4/\delta)|E(H)|$ to prove Theorem 5. Indeed, for any graph G with $|E(G)| = q \binom{m}{2} > (2/\delta)|E(H)|(m - 1)$ edges, let G' be any spanning subgraph with at least $\delta|E(G)| > 2|E(H)|(m - 1)$ edges. In this subgraph we find a copy of the matching H greedily.

When q is constant, Theorem 5 follows from an easy application of Szemerédi’s regularity lemma (see Section 3.2 for some details). Hence, we shall henceforth suppose that $q = o(1)$, $D_H \geq 2$, and (cf. (5)) $v_H \geq 2$. Now we proceed with the details of the proof.

Proof of Theorem 5. Let H be a fixed, triangle-free graph with h vertices and $e \geq 1$ edges. Without loss of generality, let δ be a constant such that $1/(\chi(H) - 1) > \delta > 0$. We start our proof by choosing the constants. Since Theorem 5 and Propositions 11, 12, and 8 involve a double alternation (“ $\forall \exists \forall \exists$ ”), this choice will consist of two rounds. In the first round (A)–(G) we address the choice of γ . After this we get the function q and we deal with the choice of M_0 in the second round (H)–(L).

- (A) Set $\alpha = \delta/16, \eta = 1/2$, and $\ell = \chi(H) > 1$.
- (B) Proposition 8 (Embedding Lemma) applied with $\alpha_{\text{EL}} = \alpha$ and $\eta_{\text{EL}} = \eta$ yields ε_{EL} and γ_{EL} .
- (C) Then we apply Proposition 12 (Pair-to-Tuple Lemma) with $d = d_H, \alpha_{\text{P2T}} = \alpha, \varepsilon_{\text{P2T}} = \varepsilon_{\text{EL}}$, and obtain δ_{P2T} and γ_{P2T} .
- (D) Proposition 11 (Regularity-to-Pair Lemma) applied with $\alpha_{\text{R2P}} = \alpha$ and $q_{\text{R2P}} = \delta_{\text{P2T}}$ yields δ_{R2P} and γ_{R2P} .
- (E) We define

$$\varepsilon = \min \left\{ \frac{\delta_{\text{R2P}}}{3}, \frac{\delta}{160} \right\}. \tag{14}$$

(F) We then apply Proposition 9 (Sparse Regularity Lemma) with $\varepsilon_{RL} = \varepsilon$, $C_{RL} = 1 + \delta/4$, and $t_{1,RL} = \max\{2\ell/\delta^{1/2}, 80/\delta\}$ and obtain $T_{1,RL} \geq t_{1,RL}$, $\xi_{RL} \leq \min\{1/2T_{1,RL}, \varepsilon\}$, and $M_{1,RL}$.

(G) Finally, we fix the constant γ promised by Theorem 5 by setting

$$\gamma = \frac{1 - \varepsilon}{T_{1,RL}} \min\{\gamma_{EL}, \gamma_{P2T}, \gamma_{R2P}, \delta\xi_{RL}/4\}. \tag{15}$$

Now let $q = q(m) = o(1)$ be a function satisfying $q \geq q_{\gamma,H}(m)$ (see Remark 6). From this we have $q^{d_H} m \gg 1$.

(H) Proposition 8 (Embedding Lemma) applied with $\alpha_{EL} = \alpha$, $\eta_{EL} = \eta$, and $p_{EL} = \alpha q$ yields N_{EL} .

(I) Then we apply Proposition 12 (Pair-to-Tuple Lemma) with $d = d_H$, $\alpha_{P2T} = \alpha$, $\varepsilon_{P2T} = \varepsilon_{EL}$, $p_{P2T} = p_{EL} = \alpha q$ and obtain N_{P2T} .

(J) Proposition 11 (Regularity-to-Pair Lemma) applied with $\alpha_{R2P} = \alpha$, $\varrho_{R2P} = \delta p_{P2T}$, and $q_{R2P} = q$ yields n_{R2P} .

(K) We use Lemma 10 (Slicing Lemma) with $\alpha_{SL} = \alpha$, $C_{SL} = 1 + \delta/4$, $\varepsilon_{SL} = \varepsilon$, and $q_{SL} = q$ to obtain n_{SL} .

(L) Finally, we let

$$M_0 = \max\left\{1 + \frac{16}{\delta^2}, M_{1,RL}, \max\{N_{EL}, N_{P2T}, n_{R2P}, n_{SL}\} \cdot T_{1,RL}\right\}.$$

Let G be any $(q, \gamma q^{v_H} m)$ -bi-jumbled graph with $m \geq M_0$ vertices, and let G' be an arbitrary spanning subgraph of G with

$$|E(G')| \geq \left(1 - \frac{1}{\ell - 1} + \delta\right) |E(G)|. \tag{16}$$

By $(q, \gamma q^{v_H} m)$ -bi-jumbledness of G and $v_H \geq 2$, we have

$$|E(G)| \geq \frac{qm^2 - \gamma q^{v_H} m^2}{2} \stackrel{(15)}{\geq} \left(1 - \frac{\delta}{4}\right) q \binom{m}{2}.$$

Hence (16) implies

$$|E(G')| \geq \left(1 - \frac{1}{\ell - 1} + \frac{3\delta}{4}\right) q \binom{m}{2}. \tag{17}$$

We claim that G' is (ξ_{RL}, C_{RL}) -bounded with respect to q . Indeed, for any two sets $X, Y \subset V(G')$, $|X|, |Y| \geq \xi_{RL} m$, we have

$$\begin{aligned} e_{G'}(X, Y) &\leq e_G(X, Y) \leq q|X||Y| + \gamma q^{v_H} m \sqrt{|X||Y|} \\ &\leq \left(1 + \gamma \sqrt{\frac{m}{|X|} \frac{m}{|Y|}}\right) q|X||Y| \leq \left(1 + \frac{\gamma}{\xi_{RL}}\right) q|X||Y| \stackrel{(15)}{\leq} C_{RL} q|X||Y|. \end{aligned}$$

Since G' has at least $M_0 \geq M_{1,RL}$ vertices, we can apply Proposition 9 (Sparse Regularity Lemma) to G' with parameters $\varepsilon_{RL} = \varepsilon$, C_{RL} , and $t_{1,RL}$ defined in (F). This yields an (ε, q) -regular (ε, G', t) -equitable partition $\bigcup_{i=0}^t V_i$ of $V(G')$ such that $t_{1,RL} \leq t \leq T_{1,RL}$, $|V_1| = \dots = |V_t| = n$, where $(1 - \varepsilon)m/t \leq n \leq m/t$, and $|V_0| \leq \varepsilon m$.

Let G_c be the subgraph of G' obtained by removing all edges in the following four sets:

$$B_1 = \{e \in E(G') : e \cap V_0 \neq \emptyset\},$$

$$B_2 = \bigcup_{1 \leq i \leq t} \{e \in E(G') : e \subset V_i\},$$

$$B_3 = \bigcup_{1 \leq i < j \leq t} \{e \in E_{G'}(V_i, V_j) : (V_i, V_j) \text{ is not } (\varepsilon, q)\text{-regular in } G'\},$$

$$B_4 = \bigcup_{1 \leq i < j \leq t} \{e \in E_{G'}(V_i, V_j) : e_{G'}(V_i, V_j) < \alpha q n^2\}.$$

Standard calculations show that

$$e(G_c) \geq \left(1 - \frac{1}{\ell - 1} + \frac{\delta}{2}\right) q \binom{m}{2} \tag{18}$$

(see [24] for details). We define the *cluster* graph F_c of G' as the graph with vertex set $V(F_c) = \{1, \dots, t\}$ and edge set

$$E(F_c) = \{\{i, j\} : (V_i, V_j) \text{ is } (\varepsilon, q)\text{-regular in } G' \text{ and } e_{G_c}(V_i, V_j) \geq \alpha q n^2\}.$$

We claim that F_c contains a copy of K_ℓ . To prove this observe first that, since G' is (ξ_{RL}, C_{RL}) -bounded with respect to q , we have $e_{G_c}(V_i, V_j) = e_{G'}(V_i, V_j) \leq C_{RL} q n^2 \leq C_{RL} q (m/t)^2$ for every $1 \leq i < j \leq t$ such that $e_{G_c}(V_i, V_j) \neq 0$.

Now using (18), the fact that $C_{RL} = 1 + \delta/4$, and the definition of F_c , we get

$$e(F_c) \geq \frac{e(G_c)}{C_{RL} q (m/t)^2} \geq \left(1 - \frac{1}{\ell - 1} + \frac{\delta}{2}\right) \left(1 - \frac{1}{m}\right) \left(1 + \frac{\delta}{4}\right)^{-1} \frac{t^2}{2}.$$

Since $m \geq M_0 > 16/\delta^2$, we have

$$e(F_c) > \left(1 - \frac{1}{\ell - 1} + \frac{\delta}{2}\right) \left(1 - \frac{\delta}{4}\right) \frac{t^2}{2} > \left(1 - \frac{1}{\ell - 1}\right) \frac{t^2}{2}.$$

The above implies that F_c contains K_ℓ as a subgraph by Turán’s theorem [34] (see also Exercise 7 on p. 189 in [7]). From this we now deduce that G' contains H as a subgraph.

Without loss of generality assume $\{1, \dots, \ell\} \subset V(F_c)$ is the vertex set of a copy of K_ℓ in F_c . Since G' is (ξ_{RL}, C_{RL}) -bounded with respect to q , it follows from the definition of F_c that the subgraphs $G_c[V_i, V_j] = G'[V_i, V_j]$, $1 \leq i < j \leq \ell$, satisfy

- (i) $|V_i| = |V_j| = n$,
- (ii) $C_{RL} q n^2 \geq e(G_c[V_i, V_j]) \geq \alpha q n^2$,
- (iii) $G_c[V_i, V_j]$ is (ε, q) -regular.

Therefore, we can apply Lemma 10 (Slicing Lemma) with $\varepsilon_{SL} = \varepsilon \leq \delta_{R2P}/3$ (see (14)), $\alpha_{SL} = \alpha$, $C_{SL} = C_{RL} = 1 + \delta/4$, and $q_{SL} = q$. This yields subgraphs $J_{ij} \subset G_c[V_i, V_j]$, $1 \leq i < j \leq \ell$, such that

- (iv) $e_{J_{ij}}(V_i, V_j) = \alpha q n^2$,
- (v) J_{ij} is $(3\varepsilon, q)$ -regular and $3\varepsilon \leq \delta_{R2P}$.

Now let Γ be the subgraph of G induced on $V_1 \cup \dots \cup V_\ell$ and let J be the subgraph of Γ defined by

$$J = \bigcup_{1 \leq i < j \leq \ell} J_{ij}.$$

Note that since G is $(q, \gamma q^{v_H} m)$ -bi-jumbled, $n \geq (1 - \varepsilon)m/t$, and $v_H \geq 2$ the graph Γ is $(q, (\gamma t/(1 - \varepsilon)\ell)q^2 \ell n)$ -bi-jumbled. It follows from (15) and $t \leq T_{1,RL}$ that $\gamma t/(1 - \varepsilon)\ell \leq \gamma_{R2P}$. Consequently, Γ is $(q, \gamma_{R2P} q^2 \ell n)$ -bi-jumbled.

Furthermore, by (iv), J is (ℓ, n, p) -partite with $p = p(n) = \alpha q(n)$. By (v), J is also (δ_{R2P}, q) -regular.

Clearly we can apply Proposition 11 (Regularity-to-Pair Lemma) with parameters $\alpha_{R2P} = \alpha$, $q_{R2P} = \delta_{P2T}$, and $q_{R2P} = q$ and deduce that J satisfies $\text{PAIR}_\ell(\delta_{P2T})$.

To conclude that J satisfies the conditions of Proposition 12 (Pair-to-Tuple Lemma), we just need to show that for all $U \subset V_i$ and $W \subset V_j$, $i \neq j \in [\ell]$, we have

$$e_J(U, W) \leq \frac{p}{\alpha} |U||W| + \gamma_{P2T} \left(\frac{p}{\alpha}\right)^{(d_H+3)/2} n \sqrt{|U||W|}.$$

From the $(q, \gamma q^{v_H} m)$ -bi-jumbledness of G , $p = \alpha q$, $v_H = (d_H + D_H + 1)/2$, and $D_H \geq 2$, we obtain

$$e_J(U, W) \leq e_G(U, W) \leq \frac{p}{\alpha} |U||W| + \gamma \left(\frac{p}{\alpha}\right)^{(D_H+d_H+1)/2} m \sqrt{|U||W|} \tag{19}$$

$$\leq \frac{p}{\alpha} |U||W| + \gamma \left(\frac{p}{\alpha}\right)^{(d_H+3)/2} m \sqrt{|U||W|}. \tag{20}$$

Hence, we just need to show that $\gamma m \leq \gamma_{P2T} n$. This, however, follows from (15), $m \leq nt/(1 - \varepsilon)$, and $t \leq T_{1,RL}$:

$$\gamma m \leq \frac{(1 - \varepsilon)\gamma_{P2T}}{T_{1,RL}} \cdot \frac{nt}{1 - \varepsilon} \leq \gamma_{P2T} n. \tag{21}$$

Thus, we can apply Proposition 12 to J with $d = d_H$, $\varepsilon_{P2T} = \varepsilon_{EL}$, and $p_{P2T} = \alpha q$ to infer that J satisfies property $\text{TUPLE}_\ell(\varepsilon_{P2T}, d_H)$.

Finally, we verify the hypothesis of Proposition 8 (Embedding Lemma). The graph J is (ℓ, n, p) -partite and has property $\text{TUPLE}_\ell(\varepsilon_{EL}, d_H)$. Similarly as in (21) we obtain $\gamma m \leq \gamma_{EL} n$. This together with (19) show that J also satisfies (12). Hence, the conditions of Proposition 8 are met and we may conclude that $J \subset G'$ contains at least $(1 - \eta_{EL})p^e n^h = p^e n^h / 2 \geq 1$ copies of H . \square

5. Sets with large neighborhoods

The motivation for the results in this section already appeared in the outline of the proof of Theorem 5. Indeed, recall that, in our discussion in Section 3.2, we defined the hypergraphs \mathcal{B}_r ($1 \leq r \leq D_H$), whose members are the r -sets B of vertices of J with the joint neighborhood $N_J(B)$ overshooting a certain bound. In what follows, we shall make this more precise and we shall prove the “local sparseness” condition of the \mathcal{B}_r mentioned in Section 3.2.

Definition 13. Let J be an (ℓ, n, p) -partite graph with ℓ -partition $\bigcup_{j=1}^{\ell} V_j$. For a given $C > 1$, we say that an s -set $S \subset V(J)$ is C -exceptionally neighborly if its common neighborhood in V_j satisfies

$$|N_J(S) \cap V_j| > C^s p^s n$$

for some $j \in [\ell]$ so that $S \cap V_j = \emptyset$. The set S is C -reasonable if it is not C -exceptionally neighborly.

The following lemma states that, under the technical hypothesis (22), in an (ℓ, n, p) -partite graph J there are only $O(\gamma^2 p^{2d-1-s} n)$ ways how to extend a C -reasonable $(s - 1)$ -set into a C -exceptionally neighborly s -set, where the implicit constant in the big O notation depends only on C, s, d , and α .

Lemma 14. For a given $0 < \alpha \leq 1$ and $d > 0$ suppose that an (ℓ, n, p) -partite graph J with ℓ -partition $\bigcup_{j=1}^{\ell} V_j$ also satisfies the property that for all $U \subset V_i$ and $W \subset V_j, i \neq j \in [\ell]$, we have

$$e_J(U, W) \leq \frac{p}{\alpha} |U||W| + \gamma \left(\frac{p}{\alpha}\right)^d n \sqrt{|U||W|}. \tag{22}$$

Let s be a positive integer, let $S \subset V(J)$ be any $(s - 1)$ -set, let $i \neq j \in [\ell]$ be such that $S \cap V_j = \emptyset$, and suppose C is such that $1/p > C > 1/\alpha \geq 1$. Set

$$W_{ij}(S) = \{w \in V_i \setminus S : |N_J(S \cup \{w\}) \cap V_j| > C^s p^s n\}.$$

If $|N_J(S) \cap V_j| \leq C^{s-1} p^{s-1} n$, then

$$|W_{ij}(S)| \leq \gamma^2 (1/\alpha)^{2d} (C - 1/\alpha)^{-2} C^{-(s-1)} p^{2d-1-s} n.$$

Proof. Let $U \subset V_j$ be a set of vertices of J containing the common neighborhood of S and of size $|U| = C^{s-1} p^{s-1} n$. Note that the definition of $W_{ij}(S)$ implies that

$$e_J(U, W_{ij}(S)) > C^s p^s n |W_{ij}(S)| = Cp|U||W_{ij}(S)|.$$

From inequality (22), we deduce that

$$(C - 1/\alpha)p|U||W_{ij}(S)| < \gamma(p/\alpha)^d n \sqrt{|U||W_{ij}(S)|},$$

whence, recalling that $|U| = C^{s-1} p^{s-1} n$, the claimed bound on $|W_{ij}(S)|$ follows. \square

We now state three corollaries that we later use in our proofs. Although these corollaries hold in more general settings, we prefer to present them in the exact way they are used later.

Corollary 15. For a given $0 < \alpha \leq 1$ and a graph H , suppose that an (ℓ, n, p) -partite graph J with ℓ -partition $\bigcup_{j=1}^{\ell} V_j$ satisfies the property that for all $U \subset V_i$ and $W \subset V_j, i \neq j \in [\ell]$, we have

$$e_J(U, W) \leq \frac{p}{\alpha} |U||W| + \gamma \left(\frac{p}{\alpha}\right)^{v_H} n \sqrt{|U||W|}.$$

Set $C = 2/\alpha > 1$ and assume $p < \alpha/2$ holds. Let S be an arbitrary subset of $V(J)$ not containing any C -exceptionally neighborly s' -set S' for every $1 \leq s' \leq D_H$. Then, for every $i \in [\ell]$, the

number of vertices $y \in V_i \setminus S$ such that $S \cup \{y\}$ contains a C -exceptionally neighborly s' -set S' for some $1 \leq s' \leq D_H$ is at most

$$2^{|S|} \ell c p^{2v_H-1-D_H} n, \tag{23}$$

where

$$c = \gamma^2 (1/\alpha)^{2v_H-2}.$$

Proof. Fix $i \in [\ell]$. Since S contains no C -exceptionally neighborly s' -set, $1 \leq s' \leq D_H$, the set $S \cup \{y\}$ ($y \in V_i \setminus S$) will contain a C -exceptionally neighborly s' -set with $1 \leq s' \leq D_H$ if and only if there exists a set of the form $S' \cup \{y\}$ with $S' \subset S$ that is C -exceptionally neighborly.

Fix $S' \subset S$ and suppose $|S'| = s' - 1$, where $1 \leq s' \leq D_H$. We apply Lemma 14, and conclude that the number of $y \in V_i \setminus S$ such that $S' \cup \{y\}$ is C -exceptionally neighborly is at most $\ell \cdot c p^{2v_H-1-s'} n$ (we multiply by ℓ to account for all possible $j \in [\ell]$). We now take the union over all $S' \subset S$, $|S'| \leq D_H - 1$, and get (23), as required. \square

Corollary 16. For a given $0 < \alpha \leq 1$ and $d > 0$ suppose that an (ℓ, n, p) -partite graph J with ℓ -partition $\bigcup_{j=1}^{\ell} V_j$ satisfies the property that for all $U \subset V_i$ and $W \subset V_j$, $i \neq j \in [\ell]$, we have

$$e_J(U, W) \leq \frac{p}{\alpha} |U||W| + \gamma \left(\frac{p}{\alpha}\right)^{(d+3)/2} n \sqrt{|U||W|}.$$

Set $C = 2/\alpha > 1$ and assume $p < \alpha/2$ holds. Then

- (a) all but at most $\ell \gamma^2 (1/\alpha)^{d+1} p^{d+1} n$ vertices $x \in V_i$ satisfy $|N_J(x) \cap V_j| \leq Cpn$ for every $j \neq i \in [\ell]$;
- (b) let $x \in V_i$ be a vertex satisfying $|N_J(x) \cap V_j| \leq Cpn$ for every $j \neq i$. Then all but at most $\ell \gamma^2 \alpha^{-d} p^d n / 2$ vertices $x' \in V_i \setminus \{x\}$ satisfy $|N_J(x, x') \cap V_j| \leq C^2 p^2 n$ for every $j \neq i$.

Outline of the proof. The first part of this corollary follows from Lemma 14 applied with $s = 1$, $S = \emptyset$, and d replaced with $(d + 3)/2$, by summing $|W_{ij}(S)|$ for all $j \in [\ell]$, $j \neq i$. In the second part we use $s = 2$ and $S = \{x\}$ instead of $S = \emptyset$. \square

6. Proof of the Embedding Lemma (Proposition 8)

The proof of Proposition 8, our Embedding Lemma, will generally follow the same lines as the proof of (*), discussed in Section 3.2. We start with some preliminary definitions and facts.

6.1. The Extension Lemma and clean embeddings

We first fix a setup under which we shall work in this section.

Setup 17. Let H and J be graphs such that

- (a) J is (ℓ, n, p) -partite with ℓ -partition $\bigcup_{j=1}^{\ell} V_j$;
- (b) H has h vertices, e edges, and an ℓ -partition $V(H) = \bigcup_{j=1}^{\ell} U_j$.

Recall that an *embedding* of H in J is an injective, edge-preserving map $f: V(H) \rightarrow V(J)$ such that $f(U_j) \subset V_j$ for all $1 \leq j \leq \ell$.

For a given $C > 1$ we say that the embedding f of H in J is (D_H, C) -reasonable if $f(H)$ contains no C -exceptionally neighborly set of size at most D_H . Denote by $\mathcal{R}(H, J; D_H, C)$ the set of all (D_H, C) -reasonable embeddings of H in J .

Moreover, for $t \in [h]$ and t -tuples $F = (u_1, \dots, u_t) \in V(H)^t$ and $X = (x_1, \dots, x_t) \in V(J)^t$, let $\mathcal{R}(H, J, F, X; D_H, C)$ denote the set of all (D_H, C) -reasonable embeddings $f \in \mathcal{R}(H, J; D_H, C)$ such that $f(u_i) = x_i$ for all $i \in [t]$. Clearly, we may always assume that all u_i , $1 \leq i \leq t$, and all x_i , $1 \leq i \leq t$, are distinct. Set $F^{\text{set}} = \{u_1, \dots, u_t\}$ and $X^{\text{set}} = \{x_1, \dots, x_t\}$.

Below, for any graph H' and any t -tuple F of vertices of H' , we write $w(H', F)$ for the number of edges in H' that do not have both endpoints in F^{set} . That is,

$$w(H', F) = |E(H')| - |E(H'[F^{\text{set}}])|.$$

Let u_1, \dots, u_h be the vertices of H . We denote by H_i , $1 \leq i \leq h$, the subgraph induced by u_1, \dots, u_i , i.e., $H_i = H[\{u_1, \dots, u_i\}]$. Recall that the ordering u_1, \dots, u_h is d -degenerate if $\deg_{H_i}(u_i) \leq d$ for all $1 \leq i \leq h$. We now state the following lemma.

Lemma 18 (*Extension Lemma*). *Let $C > 1$ be a given constant. Suppose $0 \leq t \leq \max\{2, d_H\}$, and let $F \in V(H)^t$ and $X \in V(J)^t$ be fixed. Then*

$$|\mathcal{R}(H, J, F, X; D_H, C)| \leq C^{(h-t)D_H} p^{w(H, F)} n^{h-t}.$$

In particular, if $F^{\text{set}} \subset V(H)$ is a stable set, then

$$|\mathcal{R}(H, J, F, X; D_H, C)| \leq C^{(h-t)D_H} p^e n^{h-t}.$$

Proof. It is observed in [26] that there is a D_H -degenerate ordering u_1, \dots, u_h of the vertices of H with $F^{\text{set}} = \{u_1, \dots, u_t\}$. Fix such an ordering. We shall prove

(*) for all $t \leq i \leq h$, we have

$$|\mathcal{R}(H_i, J, F, X; D_H, C)| \leq C^{(i-t)D_H} p^{w(H_i, F)} n^{i-t}, \tag{24}$$

where $H_i = H[\{u_1, \dots, u_i\}]$.

Our lemma follows from setting $i = h$ above. One may easily prove (*) by induction on i . We omit the details here (see [24]). \square

Now we derive two corollaries of Lemma 18. Denote by $\mathcal{R}_{\text{ni}}(H, J; D_H, C)$ the set of all mappings $f \in \mathcal{R}(H, J; D_H, C)$ for which $f(H)$ is a non-induced copy of H in J . The next corollary shows that the set $\mathcal{R}_{\text{ni}}(H, J; D_H, C)$ is small.

Corollary 19. *Let $C > 1$ and $\eta > 0$ be fixed and let $p = p(n) = o(1)$ be a function of n . Then there exists an integer $n_2 = n_2(p)$ such that if graphs J and H satisfy Setup 17 for $n > n_2$, then*

$$|\mathcal{R}_{\text{ni}}(H, J; D_H, C)| \leq \eta p^e n^h. \tag{25}$$

Proof. Let $\eta > 0$, $C > 1$, integers $h, \ell \geq 1$, and a function $p = p(n) = o(1)$ be given. Let $n_1 > 0$ be such that

$$p(n) \leq \frac{\eta}{h^2 C^{(h-2)(h-1)}} \tag{26}$$

for every $n > n_1$.

Suppose that graphs H and J satisfy Setup 17 with $n \geq n_1$. The case in which $h = 1$ or H is a complete graph is clear, hence we assume $h \geq 2$ and $H \neq K_h$. To count non-induced (D_H, C) -reasonable embeddings of H in J , we select an edge $\{x, x'\} \in E(J)$ and a pair u, u' of distinct, non-adjacent vertices of H . By Lemma 18 applied to $F = (u, u')$ and $X = (x, x')$, the number of (D_H, C) -reasonable embeddings $f: V(H) \rightarrow V(J)$ such that $f(u) = x$ and $f(u') = x'$ is at most $C^{(h-2)D_H} p^e n^{h-2}$.

Since $\{x, x'\} \in E(J)$ can be selected in at most pn^2 ways, the ordered pair X can be selected in at most $2pn^2$ ways. Similarly, F can be selected in at most $2\binom{h}{2}$ ways. Therefore,

$$|\mathcal{R}_{\text{ni}}(H, J; D_H, C)| \leq 4pn^2 \binom{h}{2} \cdot C^{(h-2)D_H} p^e n^{h-2} < h^2 C^{(h-2)D_H} p^{e+1} n^h.$$

The inequality $|\mathcal{R}_{\text{ni}}(H, J; D_H, C)| \leq \eta p^e n^h$ follows from $D_H \leq \Delta(H) \leq h - 1$ and (26). \square

The next two definitions introduce several important terms for our proof of Proposition 8.

Definition 20. For $\varepsilon > 0$, we call an s -set S ε -untypical if $S \cap V_j = \emptyset$ for some $j \in [\ell]$ and

$$|N_J(S) \cap V_j| \neq (1 \pm \varepsilon)p^s n.$$

To give some intuition behind Definition 21(i) below, we first recall that we are dealing with a triangle-free graph H , and hence the neighborhood of a vertex of H is stable. In view of Corollary 19, we may and shall basically disregard non-induced embeddings of H in J . Putting these two observations together, we see that we may disregard embeddings f of H in J in which we have a vertex u in $V(H)$ with $f(N_H(u))$ non-stable. Finally, we remark that, in the inductive proof that will follow, we shall be interested in avoiding ε -untypical sets for $f(N_H(u))$.

Definition 21. Let graphs J and H be as in Setup 17 and let u_1, \dots, u_h be any d_H -degenerate ordering of the vertices of H . For (i)–(iii) below, we suppose that $1 < i \leq h$.

(i) An embedding $f: V(H_{i-1}) \rightarrow V(J)$ is ε -polluted if the set $f(N_{H_i}(u_i))$ is stable but it is ε -untypical. Otherwise f is called ε -clean.

(ii) Set

$$\mathcal{R}_{\text{poll}}(H_{i-1}, J; D_H, C) = \{f \in \mathcal{R}(H_{i-1}, J; D_H, C): f \text{ is } \varepsilon\text{-polluted}\}.$$

(iii) Finally, we say that $f: V(H_{i-1}) \rightarrow V(J)$ is ε -perfect if f is ε -clean and $f(H_{i-1})$ is an induced copy of H_{i-1} in J . We also set

$$\mathcal{R}_{\text{perf}}(H_{i-1}, J; D_H, C) = \{f \in \mathcal{R}(H_{i-1}, J; D_H, C): f \text{ is } \varepsilon\text{-perfect}\}.$$

In Corollary 22 below, we estimate the size of $\mathcal{R}_{\text{poll}}(H_{i-1}, J; D_H, C)$ for $1 < i \leq h$.

Corollary 22. Let $\varepsilon > 0$ and $C > 1$ be fixed. Let J and H be graphs satisfying Setup 17 and let u_1, \dots, u_h be any d_H -degenerate ordering of the vertices of H . Suppose $1 < i \leq h$ and set $r = \deg_{H_i}(u_i)$. If J satisfies $\text{TUPLE}_\ell(\varepsilon, d_H)$ and H is triangle-free, then

$$|\mathcal{R}_{\text{poll}}(H_{i-1}, J; D_H, C)| \leq \varepsilon \ell C^{(i-1-r)D_H} p^{e(H_{i-1})} n^{i-1}.$$

In particular, if for a given $\eta > 0$ we set $\varepsilon = \varepsilon'(\eta, C, H) = \eta/\ell C^{hD_H}$, then

$$|\mathcal{R}_{\text{poll}}(H_{i-1}, J; D_H, C)| \leq \eta p^{e(H_{i-1})} n^{i-1}$$

for all $1 < i \leq h$.

Proof. By definition, an embedding $f \in \mathcal{R}(H_{i-1}, J; D_H, C)$ is ε -polluted if $f(N_{H_i}(u_i))$ is stable and ε -untypical. Fix an r -tuple F such that $F^{\text{set}} = N_{H_i}(u_i)$. Note that we have

$$\mathcal{R}_{\text{poll}}(H_{i-1}, J; D_H, C) = \bigcup_X \mathcal{R}(H_{i-1}, J, F, X; D_H, C),$$

where the union is taken over all stable and ε -untypical r -tuples X . Therefore

$$|\mathcal{R}_{\text{poll}}(H_{i-1}, J; D_H, C)| \leq \sum_X |\mathcal{R}(H_{i-1}, J, F, X; D_H, C)|, \tag{27}$$

where the sum is over the same set of r -tuples X .

Since J satisfies $\text{TUPLE}_\ell(\varepsilon, d_H)$, the number of r -tuples X that we are summing over in (27) is at most $\varepsilon \ell n^r$, where $r = \deg_{H_i}(u_i) \leq d_H$. Observe also that $N_{H_i}(u_i)$ is a stable set in H_i , because $H_i \subset H$ is triangle-free. We now apply Lemma 18 to deduce from (27) that $|\mathcal{R}_{\text{poll}}(H_{i-1}, J; D_H, C)|$ is at most

$$\varepsilon \ell n^r \cdot C^{(i-1-r)D_H} p^{e(H_{i-1})} n^{i-1-r} = \varepsilon \ell C^{(i-1-r)D_H} p^{e(H_{i-1})} n^{i-1},$$

and our corollary follows. \square

6.2. Proof of Proposition 8

Now we prove the Embedding Lemma.

Proof. Let H be any triangle-free, ℓ -partite graph with h vertices and e edges. We also fix any d_H -degenerate ordering u_1, \dots, u_h of the vertices of H , and set $H_i = H[\{u_1, \dots, u_i\}]$ for every i , $1 \leq i \leq h$.

Throughout this proof, we suppose that $0 < \alpha, \eta \leq 1$ and $C = 2/\alpha > 1$ are fixed constants. We shall prove by induction on i that

(**) for all $1 \leq i \leq h$ and all $\delta > 0$, there are $\varepsilon_i = \varepsilon_i(H, \alpha, \delta) > 0$, $\gamma_i = \gamma_i(H, \alpha, \delta) > 0$ such that for a given function $p = p(n) = o(1)$ satisfying $p^{d_H} n \gg 1$ there is $n(i) = n(i; H, \alpha, \delta, p)$ such that if

- (a) J is (ℓ, n, p) -partite and $n > n(i)$,
- (b) for all $U \subset V_j$ and $W \subset V_{j'}$, $j \neq j' \in [\ell]$, we have

$$e_J(U, W) \leq \frac{p}{\alpha} |U||W| + \gamma_i \left(\frac{p}{\alpha}\right)^{v_H} n \sqrt{|U||W|},$$

- (c) J satisfies $\text{TUPLE}_\ell(\varepsilon_i, d_H)$,
- then

$$|\mathcal{R}(H_i, J; D_H, C)| = (1 \pm \delta) p^{e(H_h)} n^i. \tag{28}$$

Note that Proposition 8 follows from (**) by taking $\delta = \eta$, $\varepsilon = \varepsilon_h$, $\gamma = \gamma_h$, and $N_1(H, \alpha, \eta, p) = n(h; H, \alpha, \eta, p)$.

Clearly, when $h \geq i = 1$, (**) holds with $\varepsilon_1 = \delta$, $\gamma_1 = n(1) = 1$ for any $\delta > 0$ and any p . Suppose now that $1 < i \leq h$ and that (**) holds for all smaller values of i .

For a given $\delta > 0$, set

$$\delta' = \min \left\{ \frac{\delta}{6}, \frac{\delta}{2\ell^{D_H} C^{D_H}} \right\}$$

and let $\varepsilon_{i-1}(H, \alpha, \delta')$ and $\gamma_{i-1}(H, \alpha, \delta')$ be given by the induction hypothesis. Furthermore, let $\varepsilon'(\delta'/2, C, H)$ be guaranteed by Corollary 22. We now set

$$\varepsilon_i = \min \left\{ \varepsilon_{i-1}(H, \alpha, \delta'), \varepsilon' \left(\frac{\delta'}{2}, C, H \right), \frac{\delta}{8} \right\}, \tag{29}$$

$$\gamma_i = \min \left\{ \left(\frac{\varepsilon_i}{2^{i-1} \ell \left(\frac{C}{2} \right)^{2\nu_H - 2}} \right)^{1/2}, \gamma_{i-1}(H, \alpha, \delta') \right\}. \tag{30}$$

For any $p = p(n) = o(1)$ with $p^{d_H} n \gg 1$, let $n(i-1) = n(i-1; H, \alpha, \delta', p)$ be given by the induction hypothesis to guarantee that

$$|\mathcal{R}(H_{i-1}, J; D_H, C)| = (1 \pm \delta') p^{e(H_{i-1})} n^{i-1} \tag{31}$$

for any graph J satisfying (a)–(c) for $n > n(i-1)$.

Now Corollary 19 tells us that for $n > n_2(\delta'/2)$,

$$|\mathcal{R}_{\text{ni}}(H_{i-1}, J; D_H, C)| \leq \frac{\delta'}{2} p^{e(H_{i-1})} n^{i-1} \tag{32}$$

holds for any (ℓ, n, p) -partite graph J .

Finally, let n_3 be such that $p^{d_H} n \geq h/\varepsilon_i$ and $p < \alpha/2$ for $n > n_3$. We set

$$n(i) = \max \{ n(i-1), n_2(\delta'/2), n_3 \}, \tag{33}$$

and claim that this choice will do. Observe that we have

$$(1 - 2\delta')(1 - 3\varepsilon_i) \geq 1 - \delta, \tag{34a}$$

$$(1 + \delta')(1 + \varepsilon_i) \leq 1 + \frac{\delta}{2}, \tag{34b}$$

$$\delta' \ell^{D_H} C^{D_H} \leq \frac{\delta}{2}. \tag{34c}$$

Let J be a graph satisfying (a)–(c) for $n > n(i)$. Then, in addition to (31) and (32), the inequality

$$|\mathcal{R}_{\text{poll}}(H_{i-1}, J; D_H, C)| \leq \frac{\delta'}{2} p^{e(H_{i-1})} n^{i-1} \tag{35}$$

also holds because $\varepsilon_i \leq \varepsilon'(\delta'/2, C, H)$ (see Corollary 22).

We start by showing the lower bound on $|\mathcal{R}(H_i, J; D_H, C)|$. Let $r = \deg_{H_i}(u_i) \leq \min\{i - 1, d_H\}$. Note that then $e(H_{i-1}) = e(H_i) - r$. By our choice of ε_i and $n(i)$, the number of embeddings in $\mathcal{R}(H_{i-1}, J; D_H, C)$ that are either ε_i -polluted or non-induced is at most

$$\frac{\delta'}{2} p^{e(H_{i-1})} n^{i-1} = \delta' p^{e(H_{i-1})} n^{i-1} = \delta' p^{e(H_i) - r} n^{i-1}$$

(see (32) and (35)). Hence, by (31), the number $|\mathcal{R}_{\text{perf}}(H_{i-1}, J; D_H, C)|$ of ε_i -perfect embeddings of H_{i-1} in J is such that

$$(1 - 2\delta') p^{e(H_i) - r} n^{i-1} < |\mathcal{R}_{\text{perf}}(H_{i-1}, J; D_H, C)| < (1 + \delta') p^{e(H_i) - r} n^{i-1}. \tag{36}$$

Given any such embedding $f' \in \mathcal{R}_{\text{perf}}(H_{i-1}, J; D_H, C)$, we estimate the number of embeddings $f \in \mathcal{R}(H_i, J; D_H, C)$ that extend f' . Let V_j be the vertex class into which we need to embed u_i .⁷ Since f' is ε_i -clean, by Definition 21 we must have that either $f'(N_{H_i}(u_i))$ is not a stable set in J , or $f'(N_{H_i}(u_i))$ is not ε_i -untypical.

Since H is triangle-free, the set $N_{H_i}(u_i)$ is a stable set in H_i . Since f' is induced, the set $f'(N_{H_i}(u_i))$ is also a stable set. Hence, the second option must be true, and, consequently,

$$|N_J(f'(N_{H_i}(u_i))) \cap V_j| - p^r n \leq \varepsilon_i p^r n. \tag{37}$$

Note that, to obtain an extension f of f' that belongs to $\mathcal{R}(H_i, J; D_H, C)$, we only need to select $f(u_i)$ in $(N_J(f'(N_{H_i}(u_i))) \cap V_j) \setminus f'(V(H_{i-1}))$ so that $f'(V(H_{i-1})) \cup \{f(u_i)\}$ does not contain a C -exceptionally neighborly s' -set for any $1 \leq s' \leq D_H$. We apply Corollary 15 with $S = f'(V(H_{i-1}))$ and obtain that at most

$$2^{i-1} \ell \gamma_i^2 \left(\frac{C}{2}\right)^{2\nu_H-2} p^{2\nu_H-1-D_H} n \stackrel{(30)}{\leq} \varepsilon_i p^r n \tag{38}$$

vertices in V_j cannot be chosen as $f(u_i)$. The last inequality follows from the fact that $2\nu_H - 1 - D_H = 2(D_H + d_H + 1)/2 - 1 - D_H = d_H \geq r$ and from (30). The reader may check that this tight inequality for the exponents of p in (38) explains why we cannot reduce the exponent of q in the hypothesis of Proposition 8 that G should be $(q, \gamma q^{\nu_H} m)$ -bi-jumbled.

From (37) it follows that the size of $(N_J(f'(N_{H_i}(u_i))) \cap V_j) \setminus f'(V(H_{i-1}))$ is at least

$$(1 - \varepsilon_i) p^r n - (h - 1) \geq (1 - 2\varepsilon_i) p^r n. \tag{39}$$

Consequently, every embedding $f' \in \mathcal{R}_{\text{perf}}(H_{i-1}, J; D_H, C)$ can be extended to an embedding $f \in \mathcal{R}(H_i, J; D_H, C)$ in at least

$$|(N_J(f'(N_{H_i}(u_i))) \cap V_j) \setminus f'(V(H_{i-1}))| - \varepsilon_i p^r n \stackrel{(39)}{>} (1 - 3\varepsilon_i) p^r n \tag{40}$$

ways. Combining (36) and (40) yields

$$|\mathcal{R}(H_i, J; D_H, C)| > (1 - 2\delta') p^{e(H_i)-r} n^{i-1} \cdot (1 - 3\varepsilon_i) p^r n \stackrel{(34a)}{\geq} (1 - \delta) p^{e(H_i)} n^i.$$

For the upper bound, we need to show that $|\mathcal{R}(H_i, J; D_H, C)| \leq (1 + \delta) p^{e(H_i)} n^i$. Fix an arbitrary $f' \in \mathcal{R}(H_{i-1}, J; D_H, C)$. The number of extensions of f' to embeddings of H_i in J is bounded from above by

$$|N_J(f'(N_{H_i}(u_i)))|. \tag{41}$$

If, furthermore, $f' \in \mathcal{R}_{\text{perf}}(H_{i-1}, J; D_H, C)$, then we know that (37) holds and hence the quantity in (41) is bounded by $(1 + \varepsilon_i) p^r n$. Combining this fact with the upper bound in (36), we obtain that the number of embeddings $f \in \mathcal{R}(H_i, J; D_H, C)$ whose restrictions to $V(H_{i-1})$ are in $\mathcal{R}_{\text{perf}}(H_{i-1}, J; D_H, C)$ is at most

$$(1 + \delta') p^{e(H_i)-r} n^{i-1} \cdot (1 + \varepsilon_i) p^r n \stackrel{(34b)}{\leq} \left(1 + \frac{\delta}{2}\right) p^{e(H_i)} n^i. \tag{42}$$

We already know that (see (31) and (36))

$$|\mathcal{R}(H_{i-1}, J; D_H, C) \setminus \mathcal{R}_{\text{perf}}(H_{i-1}, J; D_H, C)| \leq 3\delta' p^{e(H_i)-r} n^{i-1}.$$

⁷ By definition, every embedding f must preserve the vertex classes of H , that is, if $u_i \in U_j$ then $f(u_i) \in V_j$.

Since $r = \deg_J(u_i) \leq d_H \leq D_H$ and f' is (D_H, C) -reasonable, each such embedding f' gives rise to at most $C^r p^r n$ embeddings $f \in \mathcal{R}(H_i, J; D_H, C)$. Therefore, the number of embeddings $f \in \mathcal{R}(H_i, J; D_H, C)$ whose restrictions to $V(H_{i-1})$ are not in $\mathcal{R}_{\text{perf}}(H_{i-1}, J; D_H, C)$ is at most

$$3\delta' p^{e(H_i)-r} n^{i-1} \cdot C^r p^r n \stackrel{(34c)}{\leq} \frac{\delta}{2} p^{e(H_i)} n^i. \tag{43}$$

From (42) and (43) we deduce that $|\mathcal{R}(H_i, J; D_H, C)| \leq (1 + \delta)p^{e(H_i)} n^i$, as required. \square

7. Proof of the Pair-to-Tuple Lemma (Proposition 12)

Recall first the statement we are proving: for given integers $d \geq 1$ and $\ell > 1$ and reals $0 < \alpha, \varepsilon \leq 1$, we need to find $\delta > 0$ and $\gamma > 0$ such that for any function $p = p(n)$ with $p^d n \gg 1$ there exists $N_0 > 0$ with the following property: any (ℓ, n, p) -partite graph J , where $n \geq N_0$, such that

- (i) for all $U \subset V_i$ and $W \subset V_j, i \neq j \in [\ell]$, we have

$$e_J(U, W) \leq \frac{p}{\alpha} |U||W| + \gamma \left(\frac{p}{\alpha}\right)^{(d+3)/2} n \sqrt{|U||W|}, \tag{44}$$

- (ii) J possesses $\text{PAIR}_\ell(\delta)$

also satisfies $\text{TUPLE}_\ell(\varepsilon, d)$.

Let d, ℓ, α , and ε be given. Without loss of generality we may assume $d \geq 3$ because (ii) implies $\text{TUPLE}_\ell(\varepsilon, d)$ for $d = 1, 2$ and $\varepsilon \leq \delta$ (we do not need assumption (i) at all). Hence we must define δ and γ and, for a given $p = p(n)$, we must also define N_0 and then show that this choice is correct. Our proof uses a technique from [26,30] (see Lemma 26 and the proof of Lemma 43 in [26]) and is based on the following lemma, which is a well-known consequence of the Cauchy–Schwarz inequality.

Lemma 23. *For all $\varepsilon > 0$, there exists $0 < \varrho = \varrho(\varepsilon) < \varepsilon$ such that, for any family of real numbers $\{a_i \geq 0: 1 \leq i \leq M\}$ satisfying the conditions*

- (1) $\sum_{i=1}^M a_i \geq (1 - \varrho)Ma$ and
- (2) $\sum_{i=1}^M a_i^2 \leq (1 + \varrho)Ma^2$

for some $a \geq 0$, we have

$$|\{i: |a_i - a| < \varepsilon a\}| > (1 - \varepsilon)M.$$

Our application of Lemma 23 will involve the sets $\mathcal{T}(I)$ defined in Section 3.1. We first show that for any $\varrho > 0$ there is $\delta > 0$ so that if J possesses $\text{PAIR}_\ell(\delta)$, then for every fixed multiset $I = \{i_1, \dots, i_r\} \subset [\ell]$ with $3 \leq r \leq d$, and for every $j \in [\ell] \setminus I$, we can verify conditions (1) and (2) for the number $M = |\mathcal{T}(I)|$ of r -tuples $(x_1, \dots, x_r) \in \mathcal{T}(I)$, $a = p^r n$, and each a_i corresponding to $|N_{V_j}(x_1, \dots, x_r)|$ for some r -tuple $(x_1, \dots, x_r) \in \mathcal{T}(I)$. This is formally done in the next fact. In what follows we denote by

$$\sum_{(x_r) \in \mathcal{T}(I)}$$

the sum over all $(x_1, \dots, x_r) \in \mathcal{T}(I)$.

Fact 24. For every $0 < \varrho \leq 1$ there exist $\delta = \delta(d, \ell, \alpha, \varrho)$ and $\gamma = \gamma(d, \ell, \alpha, \varrho)$ such that for every $p = p(n) = o(1)$ with $p^d n \gg 1$ there is an integer $N_4 = N_4(d, \ell, \alpha, \varrho, p)$ with the following property: If an (ℓ, n, p) -partite graph J satisfies conditions (i) and (ii) above for $n \geq N_4$, then for every multiset $I = \{i_1, \dots, i_r\} \subset [\ell]$ with $3 \leq r \leq d$, and for every $j \in [\ell] \setminus I$, we have

$$(1) \quad \sum_{(x_\tau) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| = (1 \pm \varrho)n^r \cdot p^r n,$$

$$(2) \quad \sum_{(x_\tau) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)|^2 < (1 + \varrho)n^r \cdot (p^r n)^2.$$

Comparing Lemma 23 and Fact 24, one may expect that the upper bound in Fact 24(1) would not be necessary. However, it turns out that this upper bound is required in the proof of Fact 24(2).

Now we are ready to define δ and N_0 : let $\varrho = \varrho(\varepsilon^2) < \varepsilon^2$ be the constant guaranteed by Lemma 23 and $\delta = \delta(d, \ell, \alpha, \varrho/3)$, $\gamma = \gamma(d, \ell, \alpha, \varrho/3)$, and $N_4 = N_4(d, \ell, \alpha, \varrho/3, p)$ be given by Fact 24. Let $N_5 > 0$ be such that for any $n \geq N_5$ and $3 \leq r \leq d$, we have

$$\frac{n^r}{1 + \varrho/3} \leq (n - r)^r. \tag{45}$$

Set $N_0 = \max\{N_4, N_5\}$.

To prove that J satisfies $\text{TUPLE}_\ell(\varepsilon, d)$, we fix an arbitrary multiset $I = \{i_1, \dots, i_r\} \subset [\ell]$ with $3 \leq r \leq d$ and any $j \in [\ell] \setminus I$, and show that

$$|N_{V_j}(x_1, \dots, x_r)| = (1 \pm \varepsilon)p^r n$$

for all but at most εn^r r -tuples $(x_1, \dots, x_r) \in \mathcal{T}(I)$.

By (10) and (45), we have

$$\frac{n^r}{1 + \varrho/3} \leq (n - r)^r \leq M = |\mathcal{T}(I)| \leq n^r. \tag{46}$$

By our choice of constants we can apply Fact 24 and obtain

$$\begin{aligned} \sum_{(x_\tau) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| &> \left(1 - \frac{\varrho}{3}\right)n^r \cdot p^r n, \\ \sum_{(x_\tau) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)|^2 &< \left(1 + \frac{\varrho}{3}\right)n^r \cdot (p^r n)^2. \end{aligned}$$

It follows from (46) that

$$\sum_{(x_\tau) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| > (1 - \varrho)M \cdot p^r n,$$

and

$$\sum_{(x_\tau) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)|^2 < (1 + \varrho)M \cdot (p^r n)^2.$$

We are now clearly in position to apply Lemma 23 with $a = p^r n$ and $M = |\mathcal{T}(I)|$. We deduce that

$$|N_{V_j}(x_1, \dots, x_r)| = (1 \pm \varepsilon^2)p^r n = (1 \pm \varepsilon)p^r n$$

holds for at least

$$(1 - \varepsilon^2)M \stackrel{(46)}{\geq} \left(\frac{1 + \varepsilon}{1 + \varrho/3} \right) (1 - \varepsilon)n^r > (1 - \varepsilon)n^r$$

r -tuples $(x_1, \dots, x_r) \in \mathcal{T}(I)$. The last inequality follows from the fact that $\varrho/3 < \varepsilon^2 \leq 1$. What remains to be proved is Fact 24.

Proof of Fact 24. Let $0 < \varrho < 1$, in addition to $d \geq 3$, $\ell \geq 2$, and $0 < \alpha \leq 1$, be given. We first set an auxiliary constant $C = 2/\alpha > 1$ and then we let

$$\delta = \delta(d, \ell, \alpha, \varrho) = \min \left\{ \frac{\varrho}{\ell(d+2)}, \frac{\varrho}{2^{d+3}}, \frac{\varrho}{8\ell C^{2d}} \right\} \tag{47}$$

and

$$\gamma = \gamma(d, \ell, \alpha, \varrho) = \min \left\{ \sqrt{\frac{\varrho\alpha^{d+1}}{2\ell}}, \sqrt{\frac{\varrho\alpha^d}{4\ell C^d}} \right\}. \tag{48}$$

For any $p = p(n) = o(1)$, $p^d n \gg 1$, let $N_4 = N_4(d, \ell, \alpha, \varrho, p)$ be such that

$$\frac{d}{(1 - \delta)\delta} < pn, \quad \frac{1 + \varrho}{\varrho/8} < p^d n, \quad \text{and} \quad p < \frac{\alpha}{2} \tag{49}$$

for every $n \geq N_4$.

Now let an (ℓ, n, p) -partite graph J satisfies conditions (i) and (ii) for $n \geq N_4$. We fix an arbitrary multiset $I = \{i_1, \dots, i_r\} \subset [\ell]$, $3 \leq r \leq d$, and $j \in [\ell] \setminus I$.

To prove the lower bound in the first part of this fact, we observe

$$\sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| \geq \sum_{y \in V_j} \prod_{k=1}^r (|N_{V_k}(y)| - r). \tag{50}$$

Since J satisfies $\text{PAIR}_\ell(\delta)$, it follows that for all $j \in [\ell] \setminus I$ and any $i \in I$, at least $(1 - \delta)n$ vertices $y \in V_j$ satisfy

$$|N_{V_i}(y) - pn| < \delta pn. \tag{51}$$

Since I contains at most ℓ distinct numbers, there are at least $(1 - \ell\delta)n$ vertices $y \in V_j$ for which (51) holds simultaneously for all $i \in I$. Consequently, (50) yields

$$\sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| > (1 - \ell\delta)n((1 - \delta)pn - r)^r. \tag{52}$$

Applying the fact that $(a - b)^r \geq a^r - ra^{r-1}b$ for $a > b \geq 0$ and $(1 - \delta)pn > r$ by (49) to the right-hand side of (52), we obtain

$$\sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| \geq (1 - \ell\delta)^{r+1}n(pn)^r \left(1 - \frac{r}{(1 - \delta)pn} \right). \tag{53}$$

Since $r/(1 - \delta)pn \leq d/(1 - \delta)pn < \delta$ by (49), we deduce from (52) that

$$\sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| > (1 - \ell\delta)^{r+2}n^r \cdot p^r n. \tag{54}$$

Since $\delta \leq \varrho/\ell(d + 2)$ and $r \leq d$, we obtain $(1 - \ell\delta)^{r+2} \geq 1 - (r + 2)\ell\delta \geq 1 - \varrho$. Thus (54) becomes $\sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| > (1 - \varrho)n^r \cdot p^r n$.

To prove the upper bound in (1) of Fact 24, we first observe that

$$\sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| \leq \sum_{y \in V_j} \prod_{k=1}^r |N_{V_{i_k}}(y)|. \tag{55}$$

For each $y \in V_j$ such that inequality (51) is satisfied for all $i \in I$, we have

$$\prod_{k=1}^r |N_{V_{i_k}}(y)| \leq (1 + \delta)^r (pn)^r.$$

Since J satisfies $\text{PAIR}_\ell(\delta)$, there are at most $\ell\delta n$ vertices $y \in V_j$ for which inequality (51) fails for some $i \in I$.

Define the set $A = \{y \in V_j : |N_{V_i}(y)| \leq Cpn \ \forall i \in I\}$. Corollary 16(a)⁸ and (48) imply $|A| > (1 - \ell\gamma^2(1/\alpha)^{d+1}p^{d+1})n \geq (1 - (\varrho/2)p^{d+1})n$. Note that for every $y \in A$ we have

$$\prod_{k=1}^r |N_{V_{i_k}}(y)| \leq (Cpn)^r.$$

Finally, for every $y \notin A$ we have the trivial bound $\prod_{k=1}^r |N_{V_{i_k}}(y)| \leq n^r$. From (55) we obtain

$$\begin{aligned} \sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| &\leq n \cdot (1 + \delta)^r (pn)^r + \ell\delta n \cdot (Cpn)^r + \frac{\varrho}{2} p^{d+1} n \cdot n^r \\ &\stackrel{(47)}{\leq} \left(1 + \frac{\varrho}{8} + \frac{\varrho}{8} + \frac{\varrho}{2}\right) n^r \cdot p^r n < (1 + \varrho)n^r \cdot p^r n. \end{aligned}$$

This concludes the proof of the first part of Fact 24.

Now we prove the second part of Fact 24. By counting in two ways the pairs $((y_1, y_2), (x_1, \dots, x_r))$ such that $(x_1, \dots, x_r) \in \mathcal{T}(I)$ and $y_1 \neq y_2 \in N_{V_j}(x_1, \dots, x_r)$, we obtain the following inequality:

$$\sum_{(x_r) \in \mathcal{T}(I)} \binom{|N_{V_j}(x_1, \dots, x_r)|}{2} 2! \leq \sum_{(y_1, y_2) \in \mathcal{T}(\{j, j\})} \prod_{k=1}^r |N_{V_{i_k}}(y_1, y_2)|,$$

or, equivalently,

$$\begin{aligned} \sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)|^2 &\leq \sum_{(y_1, y_2) \in \mathcal{T}(\{j, j\})} \prod_{k=1}^r |N_{V_{i_k}}(y_1, y_2)| \\ &\quad + \sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)|. \end{aligned} \tag{56}$$

The second term on the right-hand side was already estimated in part (1):

$$\sum_{(x_r) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)| < (1 + \varrho)n^r p^r n \stackrel{(49)}{\leq} \frac{\varrho}{8} n^r (p^r n)^2. \tag{57}$$

⁸ Note that we need (44), $C = 2/\alpha$, and (49) to verify the assumptions of Corollary 16(a).

To estimate the first term on the right-hand side of (56), one analyzes four cases, namely:

(a) The pairs of vertices (y_1, y_2) such that

$$|N_{V_{i_k}}(y_1, y_2) - p^2n| < \delta p^2n$$

holds simultaneously for $k = 1, \dots, r$.

(b) The pairs of vertices (y_1, y_2) not included in (a) for which

$$|N_{V_{i_k}}(y_1, y_2)| \leq C^2 p^2n$$

holds simultaneously for all $k = 1, \dots, r$.

(c) The pairs of vertices (y_1, y_2) for which y_1 or $y_2 \in A$ and

$$|N_{V_{i_k}}(y_1, y_2)| > C^2 p^2n$$

holds for some $k \in \{1, \dots, r\}$.

(d) The pairs of vertices (y_1, y_2) for which $y_1 \notin A$ and $y_2 \notin A$.

One may check that the corresponding contributions to the first term on the right-hand side of (56) are bounded by

$$n^2 \cdot ((1 + \delta)p^2n)^r \stackrel{(47)}{\leq} \left(1 + \frac{\varrho}{8}\right) \cdot n^r \cdot (p^r n)^2 \quad \text{in case (a),} \tag{58}$$

$$\ell \delta n^2 \cdot (C^2 p^2n)^r \stackrel{(47)}{\leq} \frac{\varrho}{8} \cdot n^r \cdot (p^r n)^2 \quad \text{in case (b),} \tag{59}$$

$$\frac{\varrho}{4C^d} p^d n^2 \cdot (Cpn)^r \leq \frac{\varrho}{4} \cdot n^r \cdot (p^r n)^2 \quad \text{in case (c),} \tag{60}$$

$$\left(\frac{\varrho}{2} p^{d+1} n\right)^2 \cdot n^r \leq \frac{\varrho}{4} \cdot n^r \cdot (p^r n)^2 \quad \text{in case (d).} \tag{61}$$

(The calculations to derive (58)–(61) are simple; for (60), one uses Corollary 16(b). See [24] for details.) Using (56)–(61) above, we see that

$$\begin{aligned} & \sum_{(x_i) \in \mathcal{T}(I)} |N_{V_j}(x_1, \dots, x_r)|^2 \\ & \leq \left(1 + \frac{\varrho}{8} + \frac{\varrho}{8} + \frac{\varrho}{8} + \frac{\varrho}{4} + \frac{\varrho}{4}\right) n^r (p^r n)^2 < (1 + \varrho) n^r (p^r n)^2, \end{aligned}$$

which completes the proof of Fact 24(2). \square

Acknowledgments

The authors thank the referees for their helpful suggestions on the presentation of the paper.

References

[1] N. Alon, Explicit Ramsey graphs and orthonormal labelings, *Electron. J. Combin.* 1 (1994), research paper 12, approx. 8 pp. (electronic). MR 1302331 (95k:05145).
 [2] N. Alon, J. Spencer, *The Probabilistic Method*, second ed., Wiley–Intersci. Ser. Discrete Math. Optim., Wiley–Interscience [John Wiley & Sons], New York, 2000, with an appendix on the life and work of Paul Erdős. MR 2003f:60003.
 [3] L. Babai, M. Simonovits, J. Spencer, Extremal subgraphs of random graphs, *J. Graph Theory* 14 (5) (1990) 599–622. MR 1073101 (91h:05112).
 [4] F.R.K. Chung, Subgraphs of a hypercube containing no small even cycles, *J. Graph Theory* 16 (3) (1992) 273–286. MR 1168587 (93g:05073).
 [5] F.R.K. Chung, A spectral Turán theorem, *Combin. Probab. Comput.* 14 (5–6) (2005) 755–767. MR 2174654 (2006e:05108).

- [6] F.R.K. Chung, R.L. Graham, R.M. Wilson, Quasi-random graphs, *Combinatorica* 9 (4) (1989) 345–362. MR 91e:05074.
- [7] R. Diestel, *Graph Theory*, third ed., Grad. Texts in Math., vol. 173, Springer, Berlin, 2005. MR 2159259.
- [8] P. Erdős, *Some of My Favourite Unsolved Problems, A tribute to Paul Erdős*, Cambridge Univ. Press, Cambridge, 1990, pp. 467–478. MR 1117038 (92f:11003).
- [9] P. Erdős, M. Goldberg, J. Pach, J. Spencer, Cutting a graph into two dissimilar halves, *J. Graph Theory* 12 (1) (1988) 121–131. MR 928742 (89j:05043).
- [10] P. Erdős, M. Simonovits, A limit theorem in graph theory, *Studia Sci. Math. Hungar.* 1 (1966) 51–57. MR 0205876 (34 #5702).
- [11] P. Erdős, J. Spencer, Imbalances in k -colorations, *Networks* 1 (1971/1972) 379–385. MR 0299525 (45 #8573).
- [12] P. Erdős, A.H. Stone, On the structure of linear graphs, *Bull. Amer. Math. Soc.* 52 (1946) 1087–1091. MR 0018807 (8,333b).
- [13] P. Frankl, V. Rödl, Large triangle-free subgraphs in graphs without K_4 , *Graphs Combin.* 2 (2) (1986) 135–144. MR 89b:05124.
- [14] Z. Füredi, Random Ramsey graphs for the four-cycle, *Discrete Math.* 126 (1–3) (1994) 407–410. MR 94m:05131.
- [15] S. Gerke, A. Steger, The sparse regularity lemma and its applications, in: B.S. Webb (Ed.), *Surveys in Combinatorics*, Durham, 2005, in: London Math. Soc. Lecture Note Ser., vol. 327, Cambridge Univ. Press, Cambridge, 2005, pp. 227–258.
- [16] P.E. Haxell, Y. Kohayakawa, T. Łuczak, The induced size-Ramsey number of cycles, *Combin. Probab. Comput.* 4 (3) (1995) 217–239. MR 96h:05140.
- [17] S. Janson, T. Łuczak, A. Ruciński, An exponential bound for the probability of nonexistence of a specified subgraph in a random graph, in: *Random Graphs '87*, Poznań, 1987, Wiley, Chichester, 1990, pp. 73–87. MR 91m:05168.
- [18] Y. Kohayakawa, Szemerédi's regularity lemma for sparse graphs, in: *Foundations of Computational Mathematics*, Rio de Janeiro, 1997, Springer, Berlin, 1997, pp. 216–230. MR 99g:05145.
- [19] Y. Kohayakawa, B. Kreuter, A. Steger, An extremal problem for random graphs and the number of graphs with large even-girth, *Combinatorica* 18 (1) (1998) 101–120. MR 2000b:05118.
- [20] Y. Kohayakawa, T. Łuczak, V. Rödl, On K^4 -free subgraphs of random graphs, *Combinatorica* 17 (2) (1997) 173–213. MR 98h:05166.
- [21] Y. Kohayakawa, V. Rödl, Regular pairs in sparse random graphs. I, *Random Structures Algorithms* 22 (4) (2003) 359–434. MR 2004b:05187.
- [22] Y. Kohayakawa, V. Rödl, Szemerédi's regularity lemma and quasi-randomness, in: *Recent Advances in Algorithms and Combinatorics*, in: CMS Books Math./Ouvrages Math. SMC, vol. 11, Springer, New York, 2003, pp. 289–351. MR 2003j:05065.
- [23] Y. Kohayakawa, V. Rödl, M. Schacht, The Turán theorem for random graphs, *Combin. Probab. Comput.* 13 (1) (2004) 61–91. MR 2034303 (2004k:05180).
- [24] Y. Kohayakawa, V. Rödl, M. Schacht, P. Sissokho, J. Skokan, Turán's theorem for pseudo-random graphs, Tech. Report TR-2006-012-A, Emory University, 2006.
- [25] Y. Kohayakawa, V. Rödl, M. Schacht, J. Skokan, Local conditions for regularity in sparse graphs (tentative title), 2005, in preparation.
- [26] Y. Kohayakawa, V. Rödl, P. Sissokho, Embedding graphs with bounded degree in sparse pseudorandom graphs, *Israel J. Math.* 139 (2004) 93–137. MR 2041225 (2004m:05243).
- [27] J. Komlós, A. Shokoufandeh, M. Simonovits, E. Szemerédi, The regularity lemma and its applications in graph theory, in: *Theoretical Aspects of Computer Science*, Tehran, 2000, in: Lecture Notes in Comput. Sci., vol. 2292, Springer, Berlin, 2002, pp. 84–112. MR 1 966 181.
- [28] M. Krivelevich, B. Sudakov, Pseudo-random graphs, in: E. Györi, G.O.H. Katona, L. Lovász (Eds.), *More Sets, Graphs and Numbers*, in: Bolyai Soc. Math. Stud., vol. 15, Springer, Berlin/Heidelberg, 2006, pp. 199–262.
- [29] M. Krivelevich, B. Sudakov, T. Szabó, Triangle factors in sparse pseudo-random graphs, *Combinatorica* 24 (3) (2004) 403–426. MR 2085364 (2005g:05120).
- [30] T. Łuczak, P. Sissokho, personal communication.
- [31] B. Sudakov, T. Szabó, V.H. Vu, A generalization of Turán's theorem, *J. Graph Theory* 49 (2005) 187–195. MR 2145507 (2005m:05120).
- [32] E. Szemerédi, Regular partitions of graphs, in: *Problèmes combinatoires et théorie des graphes*, Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976, CNRS, Paris, 1978, pp. 399–401. MR 81i:05095.
- [33] A. Thomason, Pseudorandom graphs, in: *Random Graphs '85*, Poznań, 1985, North-Holland, Amsterdam, 1987, pp. 307–331. MR 89d:05158.
- [34] P. Turán, Eine Extremalaufgabe aus der Graphentheorie, *Mat. Fiz. Lapok* 48 (1941) 436–452. MR 0018405 (8,284j).