

Bayesian quickest detection problems for some diffusion processes

Pavel V. Gapeev*

Albert N. Shiryaev**

We study the Bayesian problems of detecting a change in the drift rate of an observable diffusion process with linear and exponential penalty costs for a detection delay. The optimal times of alarms are found as the first times at which the weighted likelihood ratios hit stochastic boundaries depending on the current observations. The proof is based on the reduction of the initial problems into appropriate three-dimensional optimal stopping problems and the analysis of the associated parabolic-type free-boundary problems. We provide closed form estimates for the value functions and the boundaries, under certain nontrivial relations between the coefficients of the observable diffusion.

1. Introduction

The problem of quickest disorder detection for an observable diffusion process seeks to determine a stopping time of alarm τ which is as close as possible to the unknown time of *disorder* (or change-point) θ at which the local drift rate of the process changes from $\mu_0(\cdot)$ to $\mu_1(\cdot)$. In the classical Bayesian formulation, it is assumed that the random time θ takes the value 0 with probability π and is exponentially distributed with parameter $\lambda > 0$ given that $\theta > 0$. An optimality criterion was proposed in [22]-[23] for the time of alarm to minimize a linear combination of the false alarm probability and the expected time delay in detecting

Mathematics Subject Classification 2000: Primary 60G40, 62M20, 34K10. Secondary 62C10, 62L15, 60J60.

Key words and phrases: Disorder detection, diffusion process, multi-dimensional optimal stopping, stochastic boundary, parabolic-type free-boundary problem, a change-of-variable formula with local time on surfaces.

* London School of Economics, Department of Mathematics, Houghton Street, London WC2A 2AE, United Kingdom; e-mail: p.v.gapeev@lse.ac.uk

** Steklov Institute of Mathematics, Russian Academy of Sciences, Gubkina Street 8, Moscow 119991, Russia; e-mail: albertsh@mi.ras.ru

the disorder correctly, for sequences of i.i.d. observations. An explicit solution of the problem of detecting a change in the constant drift rate of an observable Wiener process with the same optimality criterion was derived in [25]-[26]. The appropriate optimal stopping problem for the posterior probability of the occurrence of disorder was reduced to the associated free-boundary problem for an ordinary differential operator (see also [27; Chapter IV, Section 4] or [17; Chapter VI, Section 22]). A finite time horizon version of the Wiener disorder problem was studied in [9].

The idea of replacing the initial average time delay by a certain non-additive detection delay penalty criterion was originally introduced in [24]. The resulting Bayesian risk function was expressed through the current state of a multi-dimensional Markovian sufficient statistic having state space components which are different from the posterior probability. Such a process contained all the necessary information to determine the structure of the optimal time of alarm (see also more recent works [29], [30] and [6]). In the case of exponential penalty costs for a delay, it was observed by Poor [18] that the weighted likelihood ratio process turns out to be a one-dimensional Markovian sufficient statistic, for sequences of i.i.d. observations. This idea was taken further by Beibel [4], who solved the corresponding problem of detecting a change in the drift rate of an observable Wiener process as a generalized parking problem. Bayraktar and Dayanik [1] recognized the same property from the structure of the ordinary differential-difference equation in the free-boundary problem associated with the Bayesian problem of detecting a change in the constant intensity rate of an observable Poisson process. Some other formulations of the problem for the case of detecting a change in the arrival rate of a Poisson process, leading to the appearance of essentially multi-dimensional Markovian sufficient statistics, were studied by Bayraktar, Dayanik, and Karatzas [2]-[3]. Extensive overviews of these and other related quickest sequential change-point detection methods were provided in the monographs [28] and [19].

In the present paper, we study the Bayesian quickest disorder detection problems for observable diffusions with linear and exponential delay penalty costs. We reduce the initial problems to extended optimal stopping problems for three-dimensional Markov diffusion processes, having the posterior probability, weighted likelihood ratio, and the observations as their state space components. We show that the optimal stopping times are expressed as the first times at which the weighted likelihood ratio processes hit stochastic boundaries depending on the current state of the observation process only. We verify that the value functions and the optimal stopping boundaries are characterized by means of the associated free-boundary problem for a second-order partial differential operator. The latter turns out to be of parabolic type, because the

observation process is a one-dimensional diffusion. We also derive closed form estimates for the value functions and the boundaries for a special nontrivial subclass of observable diffusions. The Bayesian sequential testing problem for such processes was recently solved in [10]. Another related problem of transient signal detection and identification of two-sided changes in the drift rates of observable diffusion processes was considered by Pospisil, Vecer and Hadjiliadis [20].

The paper is organized as follows. In Section 2, we formulate the Bayesian quickest disorder detection problem for observable diffusion processes with linear and exponential delay penalty costs and construct the associated multi-dimensional optimal stopping problem. In Sections 3 and 4, we present the associated free-boundary problem and reduce the resulting parabolic-type partial differential operator to the normal form, which is amenable for further considerations. Applying the change-of-variable formula with local time on surfaces, obtained by Peskir [16], we verify that the solution of the free-boundary problem, which satisfies certain additional conditions, provides the solution of the initial optimal stopping problem. We derive closed form estimates for the value function and the boundary, which are uniquely determined as solutions of ordinary differential equations, under certain nontrivial relations between the coefficients of the observable diffusion. The main results are stated in Theorems 3.4 and 4.2.

2. Preliminaries

In this section, we give the Bayesian formulation of the problem (see [27; Chapter IV, Section 4] or [17; Chapter VI, Section 22] for the case of Wiener processes) in which it is assumed that one observes a sample path of the diffusion process $X = (X_t)_{t \geq 0}$ with the drift rate changing from $\mu_0(\cdot)$ to $\mu_1(\cdot)$ at some random time θ taking the value 0 with probability π and being exponentially distributed with parameter $\lambda > 0$ under $\theta > 0$.

2.1. (Formulation of the problem.) Suppose that, on a probability space $(\Omega, \mathcal{F}, P_\pi)$, there exists a standard Brownian motion $B = (B_t)_{t \geq 0}$ independent of a nonnegative random variable θ such that $P_\pi(\theta = 0) = \pi$ and $P_\pi(\theta > t | \theta > 0) = e^{-\lambda t}$, for all $t \geq 0$ and some $\lambda > 0$ fixed. Let $X = (X_t)_{t \geq 0}$ be a continuous process solving the stochastic differential equation:

$$dX_t = (\mu_0(X_t) + I(\theta \leq t)(\mu_1(X_t) - \mu_0(X_t))) dt + \sigma(X_t) dB_t \quad (2.1)$$

with $X_0 = x$, where $\mu_i(x)$, $i = 0, 1$, and $\sigma(x) > 0$ are some continuously differentiable functions on $(0, \infty)$, satisfying the conditions:

$$|\mu_i(x)| + |\sigma(x)| \leq K(1 + |x|) \quad \text{and} \quad 0 < \left| \frac{\mu_1(x) - \mu_0(x)}{\sigma(x)} \right| \leq K \quad (2.2)$$

for all $x > 0$ and some $K > 0$ fixed. In order to facilitate the considerations of the examples below, we assume the state space of the process X to be the positive half line $(0, \infty)$. It thus follows from [14; Theorem 4.6] that the equation in (2.1) admits a unique strong solution under $\theta = s$, and hence, $P_\pi(X \in \cdot | \theta = s) = P^s(X \in \cdot)$ is the distribution law of a time-homogeneous diffusion process started at some $x > 0$, with diffusion coefficient $\sigma(x)$ and the drift rate changing from $\mu_0(x)$ to $\mu_1(x)$ at time $s \in [0, \infty]$. In this case, we may conclude that the probability measure P_π has the structure:

$$P_\pi(X \in \cdot) = \pi P^0(X \in \cdot) + (1 - \pi) \int_0^\infty P^s(X \in \cdot) \lambda e^{-\lambda s} ds \quad (2.3)$$

for any $\pi \in [0, 1)$ fixed.

Based upon the continuous observation of the process X , our task is to find among the stopping times τ of X (i.e. stopping times with respect to the natural filtration $\mathcal{F}_t = \sigma(X_s | 0 \leq s \leq t)$ of the process X) an optimal time at which an *alarm* should be sounded *as close as possible* to the unobservable time of *disorder* θ . More precisely, the *Bayesian quickest detection problem* consists of computing the Bayesian risk function:

$$V(\pi) = \inf_{\tau} \left(P_\pi(\tau < \theta) + E_\pi[F(\tau - \theta)I(\tau \geq \theta)] \right) \quad (2.4)$$

and finding the optimal stopping time, called the π -Bayesian time, at which the infimum is attained in (2.4). Here $P_\pi(\tau < \theta)$ is the probability of a *false alarm*, and $E_\pi[F(\tau - \theta)I(\tau \geq \theta)]$ is the expected *costs of delay* in detecting of the disorder correctly (i.e. when $\tau \geq \theta$), where the *delay penalty function* $F(t)$ satisfies the conditions $F(t) \geq 0$ for $t \geq 0$, and $F(t) = 0$ for $t \leq 0$. We will further assume that either $F(t) = ct$ or $F(t) = c(e^{\alpha t} - 1)$ holds in (2.4) for all $t \geq 0$.

Remark 2.1. It was shown in [24], [29] and [6] that, when the Laplace transforms of delay penalty functions are of rational structure, there exist finite-dimensional processes called Markovian *sufficient statistics* in the corresponding Bayesian quickest detection problems. Such (time-homogeneous strong) Markov processes containing all the necessary information to determine the optimal stopping times (see [27; Chapter II, Section 15] for an extensive discussion of this notion). For example, the function $F(t) = ct^\delta$ for $t \geq 0$, with some $c, \delta > 0$, $\delta \in \mathbb{N}$, is of such type, while the assumption $\delta \notin \mathbb{N}$ leads to the appearance of an infinite-dimensional Markovian sufficient statistic in that case.

2.2. (Likelihood ratio and posterior probability.) In order to derive Markovian sufficient statistics for the problem of (2.4), for the cases of linear and exponential delay penalty functions

indicated above, let us define the *posterior probability* process $(\pi_t)_{t \geq 0}$ by $\pi_t = P(\theta \leq t | \mathcal{F}_t)$ for $t \geq 0$. Taking into account the fact that the probability measure P^s is equivalent to P_π on \mathcal{F}_t by construction, for any $s \in [0, \infty]$, using Bayes' formula (see, e.g. [14; Theorem 7.23]), we get that $(\pi_t)_{t \geq 0}$ admits the representation:

$$\pi_t = \pi \frac{d(P^0 | \mathcal{F}_t)}{d(P_\pi | \mathcal{F}_t)} + (1 - \pi) \int_0^t \frac{d(P^s | \mathcal{F}_t)}{d(P_\pi | \mathcal{F}_t)} \lambda e^{-\lambda s} ds. \quad (2.5)$$

Moreover, since the measure P^u coincides with P^t on \mathcal{F}_t , for all $0 \leq t \leq u$, we see that:

$$1 - \pi_t = (1 - \pi) \int_t^\infty \frac{d(P^u | \mathcal{F}_t)}{d(P_\pi | \mathcal{F}_t)} \lambda e^{-\lambda u} du = (1 - \pi) e^{-\lambda t} \frac{d(P^t | \mathcal{F}_t)}{d(P_\pi | \mathcal{F}_t)} \quad (2.6)$$

is satisfied. By means of Girsanov's theorem for diffusion processes (see, e.g. [14; Theorem 7.19]), it follows from the structure of the observation process X in (2.1) that the *likelihood ratio* process $L = (L_t)_{t \geq 0}$ defined by:

$$L_t = \frac{d(P^0 | \mathcal{F}_t)}{d(P^t | \mathcal{F}_t)} \equiv \frac{d(P^0 | \mathcal{F}_t)}{d(P^\infty | \mathcal{F}_t)} \quad (2.7)$$

admits the representation:

$$L_t = \exp \left(\int_0^t \frac{\mu_1(X_s) - \mu_0(X_s)}{\sigma^2(X_s)} dX_s - \frac{1}{2} \int_0^t \frac{\mu_1^2(X_s) - \mu_0^2(X_s)}{\sigma^2(X_s)} ds \right). \quad (2.8)$$

Hence, the expressions in (2.7) and (2.8) yield that the properties:

$$\frac{d(P^s | \mathcal{F}_t)}{d(P_\pi | \mathcal{F}_t)} \frac{d(P_\pi | \mathcal{F}_t)}{d(P^t | \mathcal{F}_t)} = \frac{d(P^s | \mathcal{F}_t)}{d(P^0 | \mathcal{F}_t)} \frac{d(P^0 | \mathcal{F}_t)}{d(P^t | \mathcal{F}_t)} = \frac{d(P^s | \mathcal{F}_s)}{d(P^0 | \mathcal{F}_s)} \frac{d(P^0 | \mathcal{F}_t)}{d(P^t | \mathcal{F}_t)} \equiv \frac{L_t}{L_s} \quad (2.9)$$

hold for each $0 \leq s \leq t$. We therefore obtain from the representations in (2.5) and (2.6) that the *weighted likelihood ratio* process $(\varphi_t)_{t \geq 0}$ defined by $\varphi_t = \pi_t / (1 - \pi_t)$ has the form:

$$\varphi_t = e^{\lambda t} L_t \left(\frac{\pi}{1 - \pi} + \int_0^t \frac{\lambda e^{-\lambda s}}{L_s} ds \right). \quad (2.10)$$

2.3. (Stochastic differential equations.) Applying Itô's formula (see, e.g. [14; Chapter IV, Theorem 4.4] or [21; Chapter IV, Theorem 3.3]) to the expression in (2.8), we get that the process L admits the representation:

$$dL_t = \frac{\mu_1(X_t) - \mu_0(X_t)}{\sigma^2(X_t)} L_t (dX_t - \mu_0(X_t) dt) \quad (2.11)$$

with $L_0 = 1$. Then, using the integration by parts formula, we see that the process $(\varphi_t)_{t \geq 0}$ from (2.10) solves the stochastic differential equation:

$$d\varphi_t = \left(\lambda(1 + \varphi_t) + \left(\frac{\mu_1(X_t) - \mu_0(X_t)}{\sigma(X_t)} \right)^2 \frac{\varphi_t^2}{1 + \varphi_t} \right) dt + \frac{\mu_1(X_t) - \mu_0(X_t)}{\sigma(X_t)} \varphi_t d\bar{B}_t \quad (2.12)$$

with $\varphi_0 = \varphi \equiv \pi/(1 - \pi)$. Hence, using Itô's formula again, we obtain that the process $(\pi_t)_{t \geq 0}$ admits the representation:

$$d\pi_t = \lambda(1 - \pi_t) dt + \frac{\mu_1(X_t) - \mu_0(X_t)}{\sigma(X_t)} \pi_t(1 - \pi_t) d\bar{B}_t \quad (2.13)$$

with $\pi_0 = \pi$. Here, the *innovation* process $\bar{B} = (\bar{B}_t)_{t \geq 0}$ defined by:

$$\bar{B}_t = \int_0^t \frac{dX_s}{\sigma(X_s)} - \int_0^t \left(\frac{\mu_0(X_s)}{\sigma(X_s)} + \pi_s \frac{\mu_1(X_s) - \mu_0(X_s)}{\sigma(X_s)} \right) ds \quad (2.14)$$

is a standard Brownian motion under the measure P_π , with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$, according to P. Lévy's characterization theorem (see, e.g. [14; Theorem 4.1] or [21; Chapter IV, Theorem 3.6]). It thus follows from (2.14) that the process X admits the representation:

$$dX_t = (\mu_0(X_t) + \pi_t(\mu_1(X_t) - \mu_0(X_t))) dt + \sigma(X_t) d\bar{B}_t \quad (2.15)$$

with $X_0 = x$. Taking into account the assumptions in (2.2), we may conclude by virtue of Remark to [14; Chapter IV, Theorem 4.6] (see also [15; Chapter V, Theorem 5.2.1]) that the processes $(\pi_t, X_t)_{t \geq 0}$ and $(\varphi_t, X_t)_{t \geq 0}$ turn out to be unique strong solutions of the corresponding systems of stochastic differential equations in (2.12), (2.13), and (2.15). According to [15; Chapter VII, Theorem 7.2.4], such processes have the (time-homogeneous strong) Markov property with respect to its natural filtration, which inherently coincides with $(\mathcal{F}_t)_{t \geq 0}$.

2.4. (Some examples.) Let us now present some expressions for the Bayesian risk functions and the appropriate Markovian sufficient statistics in the corresponding quickest disorder detection problems for observable diffusion processes.

Example 2.2. Assume that we have $F(t) = ct$ with some $c > 0$ fixed (see [25], [26], [27; Chapter IV], and [17; Chapter VI, Section 22]). It is then shown by means of standard arguments from [27; Chapter IV, Section 3] that the Bayesian risk function $V(\pi)$ in (2.4) admits the representation:

$$V'(\pi, \varphi, x) = \inf_{\tau} E_{\pi, \varphi, x} \left[1 - \pi_{\tau} + \int_0^{\tau} (1 - \pi_t) c \varphi_t dt \right] \quad (2.16)$$

where the infimum is taken over all stopping times τ such that $E_{\pi, \varphi, x} \tau < \infty$ holds. Here, $P_{\pi, \varphi, x}$ is a measure of the diffusion process $(\pi_t, \varphi_t, X_t)_{t \geq 0}$, started at some $(\pi, \varphi, x) \in [0, 1) \times [0, \infty) \times (0, \infty)$ and solving the equations in (2.12), (2.13), and (2.15), which is a Markovian sufficient statistic in the problem.

Example 2.3. Assume now that $F(t) = c(e^{\alpha t} - 1)$ with some $c, \alpha > 0$ fixed (see [24; Example 4], [18], [4], and [1]). It can be shown following the schema of arguments from [1] that the Bayesian risk function $V(\pi)$ in (2.4) admits the representation:

$$V_*(\pi, \phi, x) = \inf_{\tau} E_{\pi, \phi, x} \left[1 - \pi_{\tau} + \int_0^{\tau} (1 - \pi_t) c \alpha \phi_t dt \right] \quad (2.17)$$

where the infimum is taken over all stopping times τ such that the integral above has a finite expectation, so that $E_{\pi, \phi, x} \tau < \infty$ holds. Here, the *weighted likelihood ratio* process $(\phi_t)_{t \geq 0}$ defined by:

$$\phi_t = e^{(\alpha + \lambda)t} L_t \left(\frac{\pi}{1 - \pi} + \int_0^t \frac{\lambda e^{-(\alpha + \lambda)s}}{L_s} ds \right) \quad (2.18)$$

solves the stochastic differential equation:

$$d\phi_t = \left(\lambda + (\lambda + \alpha) \phi_t + \left(\frac{\mu_1(X_t) - \mu_0(X_t)}{\sigma(X_t)} \right)^2 \pi_t \phi_t \right) dt + \frac{\mu_1(X_t) - \mu_0(X_t)}{\sigma(X_t)} \phi_t d\bar{B}_t \quad (2.19)$$

with $\phi_0 = \phi \equiv \pi/(1 - \pi)$. In this case, $P_{\pi, \phi, x}$ is a measure of the diffusion process $(\pi_t, \phi_t, X_t)_{t \geq 0}$, started at some $(\pi, \phi, x) \in [0, 1) \times [0, \infty) \times (0, \infty)$ and solving the equations in (2.13), (2.19), and (2.15), which is a Markovian sufficient statistic in the problem.

3. The case of exponential delay penalty costs

In this section, we formulate and prove the main assertions of the paper, which are related to the quickest detection problem with exponential delay penalty costs of Example 2.3 above.

3.1. By means of the results of general theory of optimal stopping (see, e.g. [27; Chapter III] or [17; Chapter I, Section 2.1]), it follows from the structure of the reward functional in (2.17) that the optimal stopping time is given by:

$$\tau_* = \inf \{ t \geq 0 \mid V_*(\pi_t, \phi_t, X_t) = 1 - \pi_t \} \quad (3.1)$$

whenever the corresponding integral there is of finite expectation, so that $E_{\pi, \phi, x} \tau_* < \infty$ holds. In order to specify the structure of the stopping time in (3.1), we follow the arguments from [9; Subsection 2.5] and use Itô's formula to get:

$$1 - \pi_t = 1 - \pi - \int_0^t \lambda (1 - \pi_s) ds + N_t \quad (3.2)$$

where the process $N = (N_t)_{t \geq 0}$ defined by:

$$N_t = - \int_0^t \frac{\mu_1(X_s) - \mu_0(X_s)}{\sigma(X_s)} \pi_s (1 - \pi_s) d\bar{B}_s \quad (3.3)$$

is a continuous local martingale under $P_{\pi, \phi, x}$. It follows directly from (3.2) that the process $(N_{\tau \wedge t})_{t \geq 0}$ is a uniformly integrable martingale for any stopping time τ satisfying $E_{\pi, \phi, x} \tau < \infty$. Then, applying Doob's optional sampling theorem (see, e.g. [14; Theorem 3.6] or [21; Chapter II, Theorem 3.2]), we get from the expression in (3.2) that:

$$E_{\pi, \phi, x} \left[1 - \pi_\tau + \int_0^\tau (1 - \pi_t) c\alpha \phi_t dt \right] = 1 - \pi + E_{\pi, \phi, x} \int_0^\tau (1 - \pi_t) (c\alpha \phi_t - \lambda) dt \quad (3.4)$$

holds for all $(\pi, \phi, x) \in [0, 1) \times [0, \infty) \times (0, \infty)$ and any τ such that $E_{\pi, \phi, x} \tau < \infty$. Taking into account the structure of the reward in (2.17), it is seen from (3.4) that it is never optimal to stop when $\phi_t < \lambda/(c\alpha)$ for any $t \geq 0$. This shows that all the points (π, ϕ, x) such that $\phi < \lambda/(c\alpha)$ belong to the continuation region:

$$C_* = \{(\pi, \phi, x) \in [0, 1) \times [0, \infty) \times (0, \infty) \mid V_*(\pi, \phi, x) < 1 - \pi\}. \quad (3.5)$$

3.2. In order to describe the structure of the set in (3.5), let us fix some $(\pi, \phi, x) \in C_*$ and denote by $\tau_* = \tau_*(\pi, \phi, x)$ the optimal stopping time in the problem of (2.17). Then, by means of the general optimal stopping theory for Markov processes (see, e.g. [27; Chapter III] or [17; Chapter I, Section 2.2]), we conclude that:

$$V_*(\pi, \phi, x) = E_{\pi, \phi, x} \left[1 - \pi_{\tau_*} + \int_0^{\tau_*} (1 - \pi_t) c\alpha \phi_t dt \right] < 1 - \pi \quad (3.6)$$

holds. Hence, taking any ϕ' such that $\phi' < \phi$ and using the explicit expression for the process $(\phi_t)_{t \geq 0}$ through its starting point $\phi \equiv \pi/(1 - \pi)$ in (2.18), we obtain from (2.17) that the inequalities:

$$\begin{aligned} V_*(\pi, \phi', x) &\leq E_{\pi, \phi', x} \left[1 - \pi_{\tau_*} + \int_0^{\tau_*} (1 - \pi_t) c\alpha \phi_t dt \right] \\ &\leq E_{\pi, \phi, x} \left[1 - \pi_{\tau_*} + \int_0^{\tau_*} (1 - \pi_t) c\alpha \phi_t dt \right] \end{aligned} \quad (3.7)$$

are satisfied. Thus, by virtue of the inequality in (3.6), we see that $(\pi, \phi', x) \in C_*$. Taking into account the multiplicative structure of the integrand in (2.17), we can therefore extend the approach used in [18], [4], and [1], and further assume that there exists a function $g_*(x)$ such that $0 < \lambda/(c\alpha) \leq g_*(x)$ for $x > 0$, and the continuation region in (3.5) for the optimal stopping problem of (2.17) takes the form:

$$C_* = \{(\pi, \phi, x) \in [0, 1) \times [0, \infty) \times (0, \infty) \mid \phi < g_*(x)\} \quad (3.8)$$

so that the corresponding stopping region is the closure of the set:

$$D_* = \{(\pi, \phi, x) \in [0, 1) \times [0, \infty) \times (0, \infty) \mid \phi > g_*(x)\}. \quad (3.9)$$

3.3. In order to characterize the behavior of the boundary $g_*(x)$ in (3.8)-(3.9), we observe from the equation in (2.19) that the expression:

$$\int_0^{\tau_*} (1 - \pi_t) \phi_t dt = \int_0^{\tau_*} \frac{1 - \pi_t}{\lambda + \alpha} d\phi_t - \int_0^{\tau_*} \frac{1 - \pi_t}{\lambda + \alpha} (\lambda + \rho(X_t) \pi_t \phi_t) dt + N_{\tau_*}^* \quad (3.10)$$

holds for the optimal stopping time $\tau_* = \tau_*(\pi, \phi, x)$ in (2.17) such that $(\pi, \phi, x) \in C_*$. Here, the process $N^* = (N_t^*)_{t \geq 0}$ defined by:

$$N_t^* = - \int_0^t \frac{\mu_1(X_s) - \mu_0(X_s)}{\sigma(X_s)} \frac{1 - \pi_s}{\lambda + \alpha} \phi_s d\bar{B}_s \quad (3.11)$$

is a continuous local martingale under $P_{\pi, \phi, x}$, and $\rho(x)$ is the so-called *signal/noise ratio* function given by:

$$\rho(x) = \left(\frac{\mu_1(x) - \mu_0(x)}{\sigma(x)} \right)^2 \quad (3.12)$$

for any $x > 0$. Observe that the assumption that the integral in (2.17) taken up to the optimal stopping time τ_* is of finite expectation and the third inequality in (2.2) yield that the integral in the left-hand side and the second integral on the right-hand side of (3.10) are of finite expectation too. Then, taking into account the fact that $0 \leq \pi_t \leq 1$ holds for all $t \geq 0$, and assuming that the process $(N_{\tau_* \wedge t}^*)_{t \geq 0}$ is a uniformly integrable martingale under $P_{\pi, \phi, x}$ (which is the case for the process $(M_{\tau_* \wedge t})_{t \geq 0}$ from (3.37) under the conditions of Lemma 3.3 below), by means of Doob's optional sampling theorem, we get from the expression in (3.10) that:

$$E_{\pi, \phi, x} \int_0^{\tau_*} (1 - \pi_t) \phi_t dt = E_{\pi, \phi, x} \int_0^{\tau_*} \frac{1 - \pi_t}{\lambda + \alpha} d\phi_t - E_{\pi, \phi, x} \int_0^{\tau_*} \frac{1 - \pi_t}{\lambda + \alpha} (\lambda + \rho(X_t) \pi_t \phi_t) dt \quad (3.13)$$

is satisfied. Let us now take $x' > 0$ such that $x < x'$ and recall the fact that $(\pi_t, \phi_t, X_t)_{t \geq 0}$ is a time-homogeneous Markov process. Assume that $(\pi, \phi, x) \in C_*$ is chosen sufficiently close to the stopping boundary $g_*(x)$, and note that $\tau_* = \tau_*(\pi, \phi, x)$ does not depend on x' . Hence, applying the comparison results from [31] for solutions of stochastic differential equations, we obtain that the expression in (3.13) yields that:

$$\begin{aligned} V_*(\pi, \phi, x') &\leq E_{\pi, \phi, x'} \left[1 - \pi_{\tau_*} + \int_0^{\tau_*} (1 - \pi_t) c \alpha \phi_t dt \right] \\ &\leq E_{\pi, \phi, x} \left[1 - \pi_{\tau_*} + \int_0^{\tau_*} (1 - \pi_t) c \alpha \phi_t dt \right] = V_*(\pi, \phi, x) \end{aligned} \quad (3.14)$$

holds, whenever $\rho(x)$ is an increasing function on $(0, \infty)$. By virtue of the inequality in (3.14), we may therefore conclude that $(\pi, \phi, x') \in C_*$, so that the boundary $g_*(x)$ is increasing (decreasing) in (3.8)-(3.9) whenever $\rho(x)$ is increasing (decreasing) on $(0, \infty)$, respectively.

Summarizing the facts proved above, we are now ready to formulate the following assertion.

Lemma 3.1. *Suppose that $\mu_i(x)$, $i = 0, 1$, and $\sigma(x) > 0$ are continuously differentiable functions on $(0, \infty)$ in (2.1) satisfying (2.2). Assume that the process $(N_{\tau_* \wedge t}^*)_{t \geq 0}$ from (3.11) is a uniformly integrable martingale. Then the optimal Bayesian time of alarm τ_* in the quickest disorder detection problem of (2.17) has the structure:*

$$\tau_* = \inf\{t \geq 0 \mid \phi_t \geq g_*(X_t)\} \quad (3.15)$$

whenever the corresponding integral has finite expectation, so that $E_{\pi, \phi, x} \tau_* < \infty$ holds, for all $(\pi, \phi, x) \in [0, 1) \times [0, \infty) \times (0, \infty)$, and $\tau_* = 0$ otherwise. Moreover, the property:

$$g_*(x) : (0, \infty) \rightarrow (\lambda/(c\alpha), \infty) \text{ is increasing/decreasing if } \rho(x) \text{ is increasing/decreasing} \quad (3.16)$$

holds with $\rho(x)$ defined in (3.12), for all $x > 0$.

3.4. By means of standard arguments based on the application of Itô's formula, it is shown that the infinitesimal operator $\mathbb{L}_{(\pi, \phi, X)}$ of the process $(\pi_t, \phi_t, X_t)_{t \geq 0}$ from (2.13), (2.19), and (2.15) has the structure:

$$\begin{aligned} \mathbb{L}_{(\pi, \phi, X)} &= \lambda(1 - \pi) \frac{\partial}{\partial \pi} + \left(\lambda + (\lambda + \alpha) \phi + \left(\frac{\mu_1(x) - \mu_0(x)}{\sigma(x)} \right)^2 \pi \phi \right) \frac{\partial}{\partial \phi} \\ &+ (\mu_0(x) + (\mu_1(x) - \mu_0(x)) \pi) \frac{\partial}{\partial x} + (\mu_1(x) - \mu_0(x)) \left(\pi(1 - \pi) \frac{\partial^2}{\partial \pi \partial x} + \phi \frac{\partial^2}{\partial \phi \partial x} \right) \\ &+ \frac{1}{2} \left(\frac{\mu_1(x) - \mu_0(x)}{\sigma(x)} \right)^2 \left(\pi^2(1 - \pi)^2 \frac{\partial^2}{\partial \pi^2} + 2\pi(1 - \pi) \phi \frac{\partial^2}{\partial \pi \partial \phi} + \phi^2 \frac{\partial^2}{\partial \phi^2} \right) + \frac{1}{2} \sigma^2(x) \frac{\partial^2}{\partial x^2} \end{aligned} \quad (3.17)$$

for all $(\pi, \phi, x) \in [0, 1) \times [0, \infty) \times (0, \infty)$.

According to the results of the general theory of optimal stopping problems for continuous time Markov processes (see, e.g. [11], [27; Chapter III, Section 8] and [17; Chapter IV, Section 8]), we can formulate the associated *free-boundary problem* for the unknown value function $V_*(\pi, \phi, x)$ from (2.17) and the boundary $g_*(x)$ from (3.15):

$$(\mathbb{L}_{(\pi, \phi, X)} V)(\pi, \phi, x) = -(1 - \pi) c \alpha \phi \quad \text{for } (\pi, \phi, x) \in C \quad (3.18)$$

$$V(\pi, \phi, x) \Big|_{\phi=g(x)^-} = 1 - \pi \quad (\text{instantaneous stopping}) \quad (3.19)$$

$$V(\pi, \phi, x) = 1 - \pi \quad \text{for } (\pi, \phi, x) \in D \quad (3.20)$$

$$V(\pi, \phi, x) < 1 - \pi \quad \text{for } (\pi, \phi, x) \in C \quad (3.21)$$

where C and D are defined as C_* and D_* in (3.8) and (3.9) with $g(x)$ instead of $g_*(x)$, and the condition in (3.19) is satisfied for all $\pi \in [0, 1)$ and $x > 0$.

Note that the superharmonic characterization of the value function (see [7], [27; Chapter III, Section 8] and [17; Chapter IV, Section 9]) implies that $V_*(\pi, \phi, x)$ from (2.17) is the largest function satisfying (3.18)-(3.21) with the boundary $g_*(x)$.

Remark 3.2. Observe that, since the system in (3.18)-(3.21) admits multiple solutions, we need to find some additional conditions which would specify the appropriate solution providing the value function and the optimal stopping boundary for the initial problem of (2.17). In order to derive such conditions, we shall reduce the operator in (3.17) to the normal form. We also note that the fact that the stochastic differential equations for the posterior probability, the weighted likelihood ratio, and the observation process in (2.13), (2.19), and (2.15), respectively, are driven by the same (one-dimensional) innovation Brownian motion yields the property that the infinitesimal operator in (3.17) turns out to be of parabolic type.

3.5. In order to find the normal form of the operator in (3.17) and formulate the appropriate optimal stopping and free-boundary problem, we use the one-to-one correspondence transformation of processes proposed by A.N. Kolmogorov in [12]. For this, let us define the process $Y = (Y_t)_{t \geq 0}$ by:

$$Y_t = \log \phi_t - \int_z^{X_t} \frac{\mu_1(w) - \mu_0(w)}{\sigma^2(w)} dw \quad (3.22)$$

for all $t \geq 0$, and any $z > 0$ fixed. Then, taking into account the assumption that the functions $\mu_i(x)$, $i = 0, 1$, and $\sigma(x)$ are continuously differentiable on $(0, \infty)$, by means of Itô's formula, we get that the process Y admits the representation:

$$dY_t = \left(\frac{\lambda}{\phi_t} + \lambda + \alpha - \frac{\sigma^2(X_t)}{2} \left[\frac{\mu_1^2(X_t) - \mu_0^2(X_t)}{\sigma^4(X_t)} + \frac{\partial}{\partial x} \left(\frac{\mu_1(x) - \mu_0(x)}{\sigma^2(x)} \right) \Big|_{x=X_t} \right] \right) dt \quad (3.23)$$

with $Y_0 = y$ and

$$y = \log \phi - \int_z^x \frac{\mu_1(w) - \mu_0(w)}{\sigma^2(w)} dw \quad (3.24)$$

for any $z > 0$ fixed. It is seen from the equation in (3.23) that the process Y started at $y \in \mathbb{R}$ is of bounded variation. By virtue of the second inequality in (2.2), it follows from the relation in (3.22) that there exists a one-to-one correspondence between the processes (π, ϕ, X) and (π, ϕ, Y) . Hence, for any $z > 0$ fixed, the value function $V_*(\pi, \phi, x)$ from (2.17) is equal to the one of the optimal stopping problem:

$$U_*(\pi, \phi, y) = \inf_{\tau} E_{\pi, \phi, y} \left[1 - \pi_{\tau} + \int_0^{\tau} (1 - \pi_t) c \alpha \phi_t dt \right] \quad (3.25)$$

where the infimum is taken over all stopping times τ such that the integral is of finite expectation, so that $E_{\pi, \phi, y} \tau < \infty$ holds. Here, $P_{\pi, \phi, y}$ is a measure of the diffusion process $(\pi_t, \phi_t, Y_t)_{t \geq 0}$,

started at some $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$ and solving the equations in (2.10), (2.18), and (3.22). It thus follows from (3.8)-(3.9) that there exists a continuous function $h_*(y)$ such that $0 < \lambda/(c\alpha) \leq h_*(y)$ holds for $y \in \mathbb{R}$, and the optimal stopping time in the problem of (3.25) has the structure:

$$\tau_* = \inf\{t \geq 0 \mid \phi_t \geq h_*(Y_t)\} \quad (3.26)$$

whenever the corresponding integral is of finite expectation, so that $E_{\pi, \phi, y} \tau_* < \infty$ holds, and $\tau_* = 0$ otherwise.

3.6. Standard arguments then show that the infinitesimal operator $\mathbb{L}_{(\pi, \phi, Y)}$ of the process (π, ϕ, Y) from (2.13), (2.19), and (3.23) has the structure:

$$\begin{aligned} \mathbb{L}_{(\pi, \phi, Y)} &= \lambda(1 - \pi) \frac{\partial}{\partial \pi} + \left(\lambda + (\lambda + \alpha) \phi + \left(\frac{\mu_1(x(\phi, y)) - \mu_0(x(\phi, y))}{\sigma(x(\phi, y))} \right)^2 \pi \phi \right) \frac{\partial}{\partial \phi} \\ &+ \frac{1}{2} \left(\frac{\mu_1(x(\phi, y)) - \mu_0(x(\phi, y))}{\sigma(x(\phi, y))} \right)^2 \left(\pi^2(1 - \pi)^2 \frac{\partial^2}{\partial \pi^2} + 2\pi(1 - \pi) \phi \frac{\partial^2}{\partial \pi \partial \phi} + \phi^2 \frac{\partial^2}{\partial \phi^2} \right) \\ &+ \left(\frac{\lambda}{\phi} + \lambda + \alpha - \frac{\sigma^2(x(\phi, y))}{2} \left[\frac{\mu_1^2(x(\phi, y)) - \mu_0^2(x(\phi, y))}{\sigma^4(x(\phi, y))} + \frac{\partial}{\partial x} \left(\frac{\mu_1(x) - \mu_0(x)}{\sigma^2(x)} \right) \Big|_{x=x(\phi, y)} \right] \right) \frac{\partial}{\partial y} \end{aligned} \quad (3.27)$$

for all $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$. Here, because of the second inequality in (2.2), the expression for $x(\phi, y) \equiv x(\phi, y; z)$ is uniquely determined by the relation in (3.24), for any $z > 0$.

We are now ready to formulate the associated free-boundary problem for the unknown value function $U_*(\pi, \phi, y) \equiv U_*(\pi, \phi, y; z)$ from (3.25) and the boundary $h_*(y) \equiv h_*(y; z)$ from (3.26):

$$(\mathbb{L}_{(\pi, \phi, Y)} U)(\pi, \phi, y) = -(1 - \pi) c\alpha\phi \quad \text{for } \phi < h(y) \quad (3.28)$$

$$U(\pi, \phi, y) \Big|_{\phi=h(y)-} = 1 - \pi \quad (\text{instantaneous stopping}) \quad (3.29)$$

$$U(\pi, \phi, y) = 1 - \pi \quad \text{for } \phi > h(y) \quad (3.30)$$

$$U(\pi, \phi, y) < 1 - \pi \quad \text{for } \phi < h(y) \quad (3.31)$$

where the condition in (3.29) is satisfied for all $\pi \in [0, 1)$ and $y \in \mathbb{R}$. Moreover, we assume that the following conditions hold:

$$\frac{\partial U}{\partial \phi}(\pi, \phi, y) \Big|_{\phi=h(y)-} = 0 \quad (\text{smooth fit}) \quad (3.32)$$

$$\frac{\partial U}{\partial \phi}(\pi, \phi, y) \Big|_{\phi=0+} \text{ is finite} \quad (3.33)$$

and the one-sided derivative:

$$\frac{\partial U}{\partial y}(\pi, \phi, y) \Big|_{\phi=h(y)-} \text{ exists} \quad (3.34)$$

for all $\pi \in (0, 1)$, $y \in \mathbb{R}$, and any $z > 0$ fixed.

We further search for solutions of the parabolic-type free-boundary problem in (3.28)-(3.31) satisfying the conditions in (3.32)-(3.34) and such that the resulting boundaries are continuous and of bounded variation. Since such free-boundary problems cannot, in general, be solved explicitly, the existence and uniqueness of classical as well as viscosity solutions of the related variational inequalities and their connection with the optimal stopping problems have been extensively studied in the literature (see, e.g. [8], [5], [13] or [15]). It particularly follows from the results of [8; Chapter XVI, Theorem 11.1] as well as [13; Chapter V, Section 3, Theorem 14] with [13; Chapter VI, Section 4, Theorem 12] that the free-boundary problem of (3.28)-(3.31) with (3.32)-(3.34) admits a unique solution.

3.7. We continue with the following verification assertion related to the free-boundary problem in (3.28)-(3.34).

Lemma 3.3. *Suppose that $\mu_i(x)$, $i = 0, 1$, and $\sigma(x) > 0$ are continuously differentiable functions on $(0, \infty)$ in (2.1) satisfying (2.2). Assume that the function $U(\pi, \phi, y; h_*(y)) \equiv (1 - \pi)H(\phi, y; h_*(y))$ and the continuous boundary of bounded variation $h_*(y)$ form a unique solution of the free-boundary problem in (3.28)-(3.31) satisfying the conditions of (3.32)-(3.34). Then, the value function of the optimal stopping problem in (3.25) takes the form:*

$$U_*(\pi, \phi, y) = \begin{cases} (1 - \pi)H(\phi, y; h_*(y)), & \text{if } 0 \leq \phi < h_*(y) \\ 1 - \pi, & \text{if } \phi \geq h_*(y) \end{cases} \quad (3.35)$$

and $h_*(y)$ provides the optimal stopping boundary for (3.26), whenever the corresponding integral is of finite expectation, so that $E_{\pi, \phi, y} \tau_* < \infty$ holds, for all $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$.

Proof. Let us denote by $U(\pi, \phi, y)$ the right-hand side of the expression in (3.35). Hence, applying the change-of-variable formula with local time on surfaces from [16] to $U(\pi, \phi, y)$ and $h_*(y)$, and taking into account the smooth-fit condition in (3.32), we obtain:

$$U(\pi_t, \phi_t, Y_t) = U(\pi, \phi, y) + \int_0^t (\mathbb{L}_{(\pi, \phi, Y)} U)(\pi_s, \phi_s, Y_s) I(\phi_s \neq h_*(Y_s)) ds + M_t \quad (3.36)$$

where the process $M = (M_t)_{t \geq 0}$ defined by:

$$M_t = \int_0^t \frac{\partial U}{\partial \pi}(\pi_s, \phi_s, Y_s) \frac{\mu_1(X_s) - \mu_0(X_s)}{\sigma(X_s)} \pi_s (1 - \pi_s) d\bar{B}_s + \int_0^t \frac{\partial U}{\partial \phi}(\pi_s, \phi_s, Y_s) \frac{\mu_1(X_s) - \mu_0(X_s)}{\sigma(X_s)} \phi_s d\bar{B}_s \quad (3.37)$$

is a continuous local martingale under $P_{\pi, \phi, y}$ with respect to $(\mathcal{F}_t)_{t \geq 0}$.

It follows from the equation in (3.28) and the conditions of (3.30)-(3.31) that the inequality $(\mathbb{L}_{(\pi,\phi,Y)}U)(\pi,\phi,y) \geq -(1-\pi)c\alpha\phi$ holds for any $(\pi,\phi,y) \in [0,1) \times (0,\infty) \times \mathbb{R}$ such that $\phi \neq h_*(y)$, as well as $U(\pi,\phi,y) \leq 1-\pi$ is satisfied for all $(\pi,\phi,y) \in [0,1) \times (0,\infty) \times \mathbb{R}$. Recall the assumption that the boundary $h_*(y)$ is continuous and of bounded variation and the fact that the process Y from (3.22) is of bounded variation too. We thus conclude from the assumption of continuous differentiability of the functions $\mu_i(x)$, $i = 0, 1$, and $\sigma(x)$ that the time spent by the process $(\phi_t)_{t \geq 0}$ at the boundary $h_*(Y)$ is of Lebesgue measure zero, so that the indicator which appears in (3.36) can be ignored. Hence, the expression in (3.36) yields that the inequalities:

$$\begin{aligned} 1 - \pi_\tau + \int_0^\tau (1 - \pi_t) c\alpha\phi_t dt & \tag{3.38} \\ \geq U(\pi_\tau, \phi_\tau, Y_\tau) + \int_0^\tau (1 - \pi_t) c\alpha\phi_t dt & \geq U(\pi, \phi, y) + M_\tau \end{aligned}$$

hold for any stopping time τ of the process (π, ϕ, Y) started at $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$.

Let $(\tau_n)_{n \in \mathbb{N}}$ be an arbitrary localizing sequence of stopping times for the processes M . Taking the expectations with respect to the probability measure $P_{\pi,\phi,y}$ in (3.38), by means of Doob's optional sampling theorem, we get that the inequalities:

$$\begin{aligned} E_{\pi,\phi,y} \left[1 - \pi_{\tau \wedge \tau_n} + \int_0^{\tau \wedge \tau_n} (1 - \pi_t) c\alpha\phi_t dt \right] & \tag{3.39} \\ \geq E_{\pi,\phi,y} \left[U(\pi_{\tau \wedge \tau_n}, \phi_{\tau \wedge \tau_n}, Y_{\tau \wedge \tau_n}) + \int_0^{\tau \wedge \tau_n} (1 - \pi_t) c\alpha\phi_t dt \right] \\ \geq U(\pi, \phi, y) + E_{\pi,\phi,y} [M_{\tau \wedge \tau_n}] = U(\pi, \phi, y) \end{aligned}$$

hold for all $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$. Hence, letting n go to infinity and using Fatou's lemma, we obtain:

$$\begin{aligned} E_{\pi,\phi,y} \left[1 - \pi_\tau + \int_0^\tau (1 - \pi_t) c\alpha\phi_t dt \right] & \tag{3.40} \\ \geq E_{\pi,\phi,y} \left[U(\pi_\tau, \phi_\tau, Y_\tau) + \int_0^\tau (1 - \pi_t) c\alpha\phi_t dt \right] \geq U(\pi, \phi, y) \end{aligned}$$

for any stopping time τ and all $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$. By virtue of the structure of the stopping time in (3.26), it is readily seen that the inequalities in (3.40) hold with τ_* instead of τ when $\phi \geq h_*(y)$.

It remains to show that the equalities are attained in (3.40) when τ_* replaces τ , for $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$ such that $\phi < h_*(y)$. By virtue of the fact that the function $U(\pi, \phi, y)$ and the boundary $h_*(y)$ satisfy the conditions in (3.28) and (3.29), it follows

from the expression in (3.36) and the structure of the stopping time in (3.26) that the equalities:

$$U(\pi_{\tau_* \wedge \tau_n}, \phi_{\tau_* \wedge \tau_n}, Y_{\tau_* \wedge \tau_n}) + \int_0^{\tau_* \wedge \tau_n} (1 - \pi_t) c \alpha \phi_t dt = U(\pi, \phi, y) + M_{\tau_* \wedge \tau_n} \quad (3.41)$$

hold for all $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$ and any localizing sequence $(\tau_n)_{n \in \mathbb{N}}$ of M . Hence, taking into account the assumption that the integral in (2.17) taken up to τ_* is of finite expectation and using the fact that $0 \leq U(\pi, \phi, y) \leq 1$ holds, we conclude from the expression in (3.41) that the process $(M_{\tau_* \wedge t})_{t \geq 0}$ is a uniformly integrable martingale. Therefore, taking the expectations in (3.41) and letting n go to infinity, we apply the Lebesgue dominated convergence theorem to obtain the equalities:

$$\begin{aligned} & E_{\pi, \phi, y} \left[1 - \pi_{\tau_*} + \int_0^{\tau_*} (1 - \pi_t) c \alpha \phi_t dt \right] \\ &= E_{\pi, \phi, y} \left[U(\pi_{\tau_*}, \phi_{\tau_*}, Y_{\tau_*}) + \int_0^{\tau_*} (1 - \pi_t) c \alpha \phi_t dt \right] = U(\pi, \phi, y) \end{aligned} \quad (3.42)$$

for all $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$, which together with the inequalities in (3.40) directly imply the desired assertion. \square

3.8. We are now in a position to formulate the main assertion of the paper, which follows from a straightforward combination of Lemma 3.3 above and standard change-of-variable arguments. More precisely, after obtaining the solution $U_*(\pi, \phi, y) \equiv (1 - \pi)H_*(\phi, y; z)$ with $h_*(y) \equiv h_*(y; z)$ of the free-boundary problem in (3.28)-(3.31), which satisfies the conditions in (3.32)-(3.34), we put $y = y(\pi, x; z)$ and $z = x$, in order to get the solution of the initial quickest detection problem with exponential penalty costs for a detection delay stated in (2.17).

Theorem 3.4. *Suppose that the assumptions of Lemmas 3.1 and 3.3 hold. Then, in the quickest disorder detection problem of (2.17) for the observation process X from (2.1), the Bayesian risk function takes the form $V_*(\pi, \phi, x) = U_*(\pi, \phi, y(\phi, x)) \equiv (1 - \pi)H_*(\phi, y(\phi, x; x); x)$ and the optimal stopping boundary $0 < \lambda/(c\alpha) \leq g_*(x)$ in (3.15) satisfying (3.16) is uniquely determined by the equation $g(x) = h_*(y(g(x), x)) \equiv h_*(y(g(x), x; x); x)$, for each $x > 0$ fixed. Here the function $U_*(\pi, \phi, y) \equiv (1 - \pi)H_*(\phi, y; z)$ and the continuous boundary of bounded variation $h_*(y) \equiv h_*(y; z)$ form a unique solution of the free-boundary problem in (3.28)-(3.34), and the expression for $y(\phi, x) \equiv y(\phi, x; z)$ is explicitly determined by the relation in (3.24), for all $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$ and any $z > 0$ fixed.*

Remark 3.5. Observe that the optimal stopping time in the problem of (2.17) does not depend on the dynamics of the process $(\pi_t)_{t \geq 0}$, so that the two-dimensional process $(\phi_t, X_t)_{t \geq 0}$ turns out to be a sufficient statistic. This fact is recognized as a consequence of the structure

of the partial differential equation in (3.17)-(3.18). However, the process $(\phi_t, X_t)_{t \geq 0}$ is not Markovian, and thus, in order to solve the optimal stopping problem of (2.17), we need to add the component $(\pi_t)_{t \geq 0}$ and then operate with the resulting Markov process $(\pi_t, \phi_t, X_t)_{t \geq 0}$.

Let us now give a short note concerning the case of bounded signal/noise ratio function $\rho(x)$ from (3.12).

Remark 3.6. Suppose that there exist some $0 < \underline{\rho} < \bar{\rho} < \infty$ such that $\underline{\rho} \leq \rho(x) \leq \bar{\rho}$ holds for all $x > 0$. Let us denote by $\underline{V}_*(\pi, \phi, x)$ with $\underline{g}_*(x)$ and by $\bar{V}_*(\pi, \phi, x)$ with $\bar{g}_*(x)$ the solution of the Bayesian quickest disorder detection problem with exponential delay penalty, under $\rho(x) \equiv \underline{\rho}$ and $\rho(x) \equiv \bar{\rho}$, respectively. In those cases, the problem of (2.17) degenerates into an optimal stopping problem for the two-dimensional Markov process $(\pi_t, \phi_t)_{t \geq 0}$, and the value functions $\underline{V}_*(\pi, \phi, x) \equiv \underline{V}_*(\pi, \phi)$ and $\bar{V}_*(\pi, \phi, x) \equiv \bar{V}_*(\pi, \phi)$ with the stopping boundaries $\underline{g}_*(x) \equiv \underline{h}_*$ and $\bar{g}_*(x) \equiv \bar{h}_*$ are given by the expressions in (3.55) and (3.54) below, whenever $\eta = 1/\underline{\rho}$ and $\eta = 1/\bar{\rho}$, respectively. Taking into account the properties of the boundary $g_*(x)$ in (3.16) and the fact that $V_*(\pi, \phi, x) = 1 - \pi$ for all $\phi \geq g_*(x)$ and $0 \leq \pi < 1$, we therefore conclude by standard comparison arguments that the inequalities $\bar{V}_*(\pi, \phi) \leq V_*(\pi, \phi, x) \leq \underline{V}_*(\pi, \phi)$ and thus $0 < \lambda/(c\alpha) \leq \underline{h}_* \leq g_*(x) \leq \bar{h}_*$ hold for all $(\pi, \phi, x) \in [0, 1) \times [0, \infty) \times (0, \infty)$.

3.9. In order to pick up some special cases in which the free-boundary problem in (3.28)-(3.34) can admit a simpler structure, for the rest of the section, in addition to the conditions in (2.2), we suppose that the property:

$$\mu_i(x) = \frac{\eta_i \sigma^2(x)}{x} \quad \text{for some } \eta_i \in \mathbb{R}, i = 0, 1, \quad \text{such that } \eta_0 \neq \eta_1 \quad \text{and} \quad \eta_0 + \eta_1 = 1 \quad (3.43)$$

holds for all $x > 0$. Moreover, we assume that the diffusion coefficient $\sigma(x)$ satisfies:

$$\sigma(x) \sim A_0 x^\alpha \quad \text{as } x \downarrow 0 \quad \text{and} \quad \sigma(x) \sim A_\infty x^\beta \quad \text{as } x \uparrow \infty \quad (3.44)$$

with some $A_0, A_\infty > 0$ and $\alpha, \beta \in \mathbb{R}$ such that $(1 - \alpha)\eta \leq 0$ and $(1 - \beta)\eta \geq 0$ holds, where we set $\eta = 1/(\eta_1 - \eta_0)$. In this case, the process $Y = (Y_t)_{t \geq 0}$ takes the form:

$$Y_t = \log \phi_t - \frac{1}{\eta} \log \frac{X_t}{z} \equiv \log \phi + \int_0^t \left(\frac{\lambda}{\phi_s} + \lambda + \alpha \right) ds \quad \text{with} \quad \eta = \frac{1}{\eta_1 - \eta_0} \quad (3.45)$$

for any $z > 0$ fixed. It is easily seen from the structure of the expression in (3.45) that the one-to-one correspondence between the processes $(\pi_t, \phi_t, X_t)_{t \geq 0}$ and $(\pi_t, \phi_t, Y_t)_{t \geq 0}$ remains true in this case. Hence, getting the expression for X_t from (3.45) and substituting it into the equations of (2.13) and (2.19), we obtain:

$$d\pi_t = \lambda(1 - \pi_t) dt + \frac{\sigma(z e^{-\eta Y_t} \phi_t^\eta)}{\eta z e^{-\eta Y_t} \phi_t^\eta} \pi_t (1 - \pi_t) d\bar{B}_t \quad (3.46)$$

with $\pi_0 = \pi$ and

$$d\phi_t = \left(\lambda + (\lambda + \alpha)\phi_t + \frac{\sigma^2(ze^{-\eta Y_t}\phi_t^\eta)}{\eta^2 z^2 e^{-2\eta Y_t}\phi_t^{2\eta}} \pi_t \phi_t \right) dt + \frac{\sigma(ze^{-\eta Y_t}\phi_t^\eta)}{\eta z e^{-\eta Y_t}\phi_t^\eta} \phi_t d\bar{B}_t \quad (3.47)$$

with $\phi_0 = \phi$, for any $z > 0$ fixed. Applying Itô's formula to the expression in (3.45) and taking into account the representations in (2.13) and (2.15) as well as the assumption of (3.43), we get:

$$dY_t = \left(\frac{\lambda}{\phi_t} + \lambda + \alpha \right) dt \quad (3.48)$$

with $Y_0 = y$. It thus follows that the infinitesimal operator $\mathbb{L}_{(\pi, \phi, Y)}$ from (3.27) takes the form:

$$\begin{aligned} \mathbb{L}_{(\pi, \phi, Y)} &= \lambda(1 - \pi) \frac{\partial}{\partial \pi} + \left(\lambda + (\lambda + \alpha)\phi + \frac{\sigma^2(ze^{-\eta y}\phi^\eta)}{\eta^2 z^2 e^{-2\eta y}\phi^{2\eta}} \pi \phi \right) \frac{\partial}{\partial \phi} \\ &+ \frac{1}{2} \frac{\sigma^2(ze^{-\eta y}\phi^\eta)}{\eta^2 z^2 e^{-2\eta y}\phi^{2\eta}} \left(\pi^2(1 - \pi)^2 \frac{\partial^2}{\partial \pi^2} + 2\pi(1 - \pi)\phi \frac{\partial^2}{\partial \pi \partial \phi} + \phi^2 \frac{\partial^2}{\partial \phi^2} \right) + \left(\frac{\lambda}{\phi} + \lambda + \alpha \right) \frac{\partial}{\partial y} \end{aligned} \quad (3.49)$$

for all $(\pi, \phi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$ and any $z > 0$ fixed.

3.10. Let us now introduce the function $\hat{U}(\pi, \phi, y) \equiv (1 - \pi)\hat{H}(\phi, y)$ and the boundary $\hat{h}(y)$ as a solution of the free-boundary problem consisting of the differential equation:

$$\left((\lambda + (\lambda + \alpha)\phi) \frac{\partial H}{\partial \phi} + \frac{1}{2} \frac{\sigma^2(ze^{-\eta y}\phi^\eta)}{\eta^2 z^2 e^{-2\eta y}\phi^{2\eta}} \phi^2 \frac{\partial^2 H}{\partial \phi^2} - \lambda H \right) (\phi, y) = -c\alpha\phi \quad \text{for } \phi < h(y) \quad (3.50)$$

instead of the one in (3.28), for each $y > 0$ fixed, and the conditions of (3.29)-(3.31) as well as (3.32)-(3.34). The general solution of the resulting second-order *ordinary* differential equation in (3.50) takes the form:

$$H(\phi, y) = C_0(y) H_0(\phi, y) + C_\infty(y) H_\infty(\phi, y) - c(1 + \phi) \quad (3.51)$$

where $H_i(\phi, y)$, $i = 0, \infty$, form a system of fundamental positive solutions (i.e. nontrivial linearly independent particular solutions) of the corresponding *homogeneous* differential equation, and $C_i(y)$, $i = 0, \infty$, are some arbitrary continuously differentiable functions, so that the condition in (3.34) holds. By virtue of the assumptions of (2.2) and taking into account the arguments from [10; Section 4], we can identify by $H_0(\phi, y)$ a decreasing solution that has a singularity at zero and by $H_\infty(\phi, y)$ an increasing solution that has a singularity at infinity.

Observe that we should have $C_0(y) = 0$ in (3.51), since otherwise $U(\pi, \phi, y) \equiv (1 - \pi)H(\phi, y) \rightarrow \pm\infty$ as $\phi \downarrow 0$, that must be excluded by virtue of the obvious fact that the value function in (3.25) is bounded at $\phi = 0$, for any $y \in \mathbb{R}$ fixed. Then, applying the conditions of (3.29) and (3.32) to the function in (3.51) with $C_0(y) = 0$, we get that the equalities:

$$C_\infty(y) H_\infty(h(y), y) = c(1 + h(y)) + 1 \quad \text{and} \quad C_\infty(y) \frac{\partial H_\infty}{\partial \phi}(\phi, y) \Big|_{\phi=h(y)} = c \quad (3.52)$$

hold for $y \in \mathbb{R}$ fixed. Hence, solving the equations of (3.52), we get that the solution of the system of (3.50) with (3.29) and (3.32)-(3.33) is given by:

$$H(\phi, y; \widehat{h}(y)) = (c(1 + \widehat{h}(y)) + 1) \frac{H_\infty(\phi, y)}{H_\infty(\widehat{h}(y), y)} - c(1 + \phi) \quad (3.53)$$

for all $0 \leq \phi < \widehat{h}(y)$, so that $0 \leq H(\phi, y; \widehat{h}(y)) \equiv H(\phi, y; z; \widehat{h}(y; z)) \leq 1$ holds, where $\widehat{h}(y)$ satisfies the equation:

$$\left. \frac{\partial H_\infty}{\partial \phi}(\phi, y) \right|_{\phi=h(y)} = \frac{cH_\infty(h(y), y)}{c(1 + h(y)) + 1} \quad (3.54)$$

for any $y \in \mathbb{R}$ fixed.

Taking into account the facts proved above, let us formulate the following assertion.

Corollary 3.7. *Suppose that $\mu_i(x)$, $i = 0, 1$, and $\sigma(x) > 0$ are continuously differentiable functions on $(0, \infty)$ in (2.1) satisfying (2.2) and (3.43)-(3.44) with $\alpha, \beta \in \mathbb{R}$ such that $(1 - \alpha)\eta \leq 0$ and $(1 - \beta)\eta \geq 0$, where $\eta = 1/(\eta_1 - \eta_0)$. Assume that $\widehat{h}(y)$ provides a unique solution of the equation in (3.54) for all $y \in \mathbb{R}$. Then, using the same arguments as in the proof of Lemma 3.3 above, it is shown that the function:*

$$\widehat{U}(\pi, \phi, y) \equiv (1 - \pi)\widehat{H}(\phi, y) \quad \text{with} \quad \widehat{H}(\phi, y) = \begin{cases} H(\phi, y; \widehat{h}(y)), & \text{if } 0 \leq \phi < \widehat{h}(y) \\ 1, & \text{if } \phi \geq \widehat{h}(y) \end{cases} \quad (3.55)$$

coincides with the value function of the optimal stopping problem:

$$\begin{aligned} & \widehat{U}(\pi, \phi, y) \\ &= \inf_{\tau} E_{\pi, \phi, y} \left[1 - \pi_\tau + \int_0^\tau (1 - \pi_t) \left(c\alpha\phi_t - \left(\frac{\lambda}{\phi_t} + \lambda + \alpha \right) \frac{\partial \widehat{H}}{\partial y}(\phi_t, Y_t) I(\phi_t < \widehat{h}(Y_t)) \right) dt \right] \end{aligned} \quad (3.56)$$

which corresponds to the Bayesian risk function in (3.25). Moreover, $\widehat{h}(y) \equiv \widehat{h}(y; z)$ determined by (3.54) provides a hitting boundary for the stopping time:

$$\widehat{\tau} = \inf\{t \geq 0 \mid \phi_t \geq \widehat{h}(Y_t)\} \quad (3.57)$$

which turns out to be optimal in (3.56) whenever the integral above is of finite expectation, and $\widehat{\tau} = 0$ otherwise, for any $z > 0$ fixed.

Remark 3.8. Note that the function $\widehat{U}(\pi, \phi, y)$ in (3.56) and the boundary $\widehat{h}(y)$ in (3.57) provide lower (upper) and upper (lower) estimates for the initial value function $U_*(\pi, \phi, y)$ in (3.25) and the optimal stopping boundary $h_*(y)$ in (3.26), whenever the function $y \mapsto \widehat{H}(\phi, y)$ is increasing (decreasing) on \mathbb{R} . According to Remark 3.6 above and the structure of the change of variables in (3.24), such a situation occurs when $\rho(x)$ from (3.12) is an increasing (decreasing) function on $(0, \infty)$ and $\eta_0 < \eta_1$ ($\eta_0 > \eta_1$) in (3.43), respectively.

4. The case of linear delay penalty costs

In this section, we provide some results, which are related to the quickest detection problem with linear delay penalty costs of Example 2.2 above.

4.1. Following the arguments of Subsection 3.1 above and applying Doob's optional sampling theorem, we get from (3.2) that the equality:

$$E_{\pi, \varphi, x} \left[1 - \pi_\tau + \int_0^\tau (1 - \pi_t) c \varphi_t dt \right] = 1 - \pi + E_{\pi, \varphi, x} \int_0^\tau (1 - \pi_t) (c \varphi_t - \lambda) dt \quad (4.1)$$

holds for all $(\pi, \varphi, x) \in [0, 1) \times [0, \infty) \times (0, \infty)$ and any stopping time τ satisfying $E_{\pi, \varphi, x} < \infty$. Taking into account the structure of the reward in (2.16), it is also seen from (4.1) that it is never optimal to stop when $\varphi_t < \lambda/c$ for any $t \geq 0$. This shows that all the points (π, φ, x) such that $\varphi < \lambda/c$ belong to the continuation region:

$$C' = \{(\pi, \varphi, x) \in [0, 1) \times [0, \infty) \times (0, \infty) \mid V'(\pi, \varphi, x) < 1 - \pi\}. \quad (4.2)$$

Then, combining the arguments in [27; Chapter IV, Section 3] with the ones in Subsection 3.2 above, we obtain that the continuation region in (4.2) for the optimal stopping problem of (2.16) takes the form:

$$C' = \{(\pi, \varphi, x) \in [0, 1) \times [0, \infty) \times (0, \infty) \mid \varphi < g'(x)\} \quad (4.3)$$

so that the corresponding stopping region is the closure of the set:

$$D' = \{(\pi, \varphi, x) \in [0, 1) \times [0, \infty) \times (0, \infty) \mid \varphi > g'(x)\}. \quad (4.4)$$

4.2. In order to characterize the behavior of the boundary $g'(x)$ in (4.3)-(4.4), we observe from the equation in (2.12) that the expression:

$$\int_0^{\tau'} (1 - \pi_t) \varphi_t dt = \int_0^{\tau'} \frac{1 - \pi_t}{\lambda} d\varphi_t - \int_0^{\tau'} \frac{1 - \pi_t}{\lambda} (\lambda + \rho(X_t) \pi_t \varphi_t) dt + N'_{\tau'} \quad (4.5)$$

holds for the optimal stopping time $\tau' = \tau'(\pi, \varphi, x)$ in (2.16) such that $(\pi, \varphi, x) \in C'$. Here the process $N' = (N'_t)_{t \geq 0}$ defined by:

$$N'_t = - \int_0^t \frac{\mu_1(X_s) - \mu_0(X_s)}{\sigma(X_s)} \frac{1 - \pi_s}{\lambda} \varphi_s d\bar{B}_s \quad (4.6)$$

is a continuous local martingale under $P_{\pi, \varphi, x}$, and the function $\rho(x)$ is given by (3.12). Note that the assumption that $E_{\pi, \varphi, x} \tau' < \infty$ holds and the third inequality in (2.2) yield that the integral in the left-hand side and the second integral on the right-hand side of (4.5) are of finite

expectation. Moreover, by virtue of the facts that $(1 - \pi_t)\varphi_t = \pi_t$ and $0 \leq \pi_t \leq 1$ holds for all $t \geq 0$, and taking into account the third inequality in (2.2), we see from (4.6) that the process $(N'_{\tau' \wedge t})_{t \geq 0}$ is a uniformly integrable martingale under $P_{\pi, \varphi, x}$. Then, applying Doob's optional sampling theorem, we get from the expression in (4.5) that:

$$E_{\pi, \varphi, x} \int_0^{\tau'} (1 - \pi_t) \varphi_t dt = E_{\pi, \varphi, x} \int_0^{\tau'} \frac{1 - \pi_t}{\lambda} d\varphi_t - E_{\pi, \varphi, x} \int_0^{\tau'} \frac{1 - \pi_t}{\lambda} (\lambda + \rho(X_t) \pi_t \varphi_t) dt \quad (4.7)$$

is satisfied. Let us now take $x' > 0$ such that $x < x'$ and recall the fact that $(\pi_t, \varphi_t, X_t)_{t \geq 0}$ is a time-homogeneous Markov process. Assume that $(\pi, \varphi, x) \in C'$ is chosen sufficiently close to the stopping boundary $g'(x)$, and note that $\tau' = \tau'(\pi, \varphi, x)$ does not depend on x' . Hence, by means of the comparison results for solutions of stochastic differential equations, we obtain that the expression in (4.7) yields:

$$\begin{aligned} V'(\pi, \varphi, x') - (1 - \pi) &\leq E_{\pi, \varphi, x'} \left[1 - \pi_{\tau_*} + \int_0^{\tau_*} (1 - \pi_t) c \varphi_t dt \right] \\ &\leq E_{\pi, \varphi, x} \left[1 - \pi_{\tau_*} + \int_0^{\tau_*} (1 - \pi_t) c \varphi_t dt \right] = V'(\pi, \varphi, x) - (1 - \pi) \end{aligned} \quad (4.8)$$

whenever $\rho(x)$ is an increasing function on $(0, \infty)$. By virtue of the inequality in (4.8), we may therefore conclude that $(\pi, \varphi, x') \in C'$, so that the boundary $g'(x)$ is increasing (decreasing) in (4.3)-(4.4) whenever $\rho(x)$ is increasing (decreasing) on $(0, \infty)$, respectively.

Summarizing the facts proved above, we now formulate the assertions related to the Bayesian quickest detection problem with linear penalty costs for a detection delay, which are proved using the arguments from the previous section.

Lemma 4.1. *Suppose that $\mu_i(x)$, $i = 0, 1$, and $\sigma(x) > 0$ are continuously differentiable functions on $(0, \infty)$ in (2.1) satisfying (2.2). Then the optimal Bayesian time of alarm τ' in the quickest disorder detection problem (2.16) has the structure:*

$$\tau' = \inf\{t \geq 0 \mid \varphi_t \geq g'(X_t)\} \quad (4.9)$$

whenever $E_{\pi, \varphi, x} \tau' < \infty$ holds, for all $(\pi, \varphi, x) \in [0, 1) \times [0, \infty) \times (0, \infty)$, and $\tau' = 0$ otherwise. Moreover, the property:

$$g'(x) : (0, \infty) \rightarrow (\lambda/c, \infty) \text{ is increasing/decreasing if } \rho(x) \text{ is increasing/decreasing} \quad (4.10)$$

holds with $\rho(x)$ defined in (3.12), for all $x > 0$.

Theorem 4.2. *Suppose that the assumptions of Lemmas 4.1 and 3.3 hold with $\phi = \varphi$, $\alpha = 0$ in (3.27), and $\alpha = 1$ in (3.28). Then, in the quickest disorder detection problem of*

(2.16) for the observation process X from (2.1), the Bayesian risk function takes the form $V'(\pi, \varphi, x) = U'(\pi, \varphi, y(\varphi, x)) \equiv (1 - \pi)H'(\varphi, y(\varphi, x); x)$ and the optimal stopping boundary $0 < \lambda/c \leq g'(x)$ in (4.9) satisfying (4.10) is uniquely determined by the equation $g(x) = h'(y(g(x), x)) \equiv h'(y(g(x), x); x)$, for each $x > 0$ fixed. Here the function $U'(\pi, \varphi, y) \equiv (1 - \pi)H'(\varphi, y; z)$ and the bounded continuous boundary of bounded variation $h'(y) \equiv h'(y; z)$ form a unique solution of the free-boundary problem in (3.28)-(3.34), and the expression for $y(\varphi, x) \equiv y(\varphi, x; z)$ is explicitly determined by the relation in (3.24) with $\phi = \varphi$, for all $(\pi, \varphi, y) \in [0, 1) \times (0, \infty) \times \mathbb{R}$ and any $z > 0$ fixed.

Remark 4.3. Suppose that there exist some $0 < \underline{\rho} < \bar{\rho} < \infty$ such that $\underline{\rho} \leq \rho(x) \leq \bar{\rho}$ holds for all $x > 0$. Let us denote by $\underline{V}'(\pi, \varphi, x)$ with $\underline{g}'(x)$ and by $\bar{V}'(\pi, \varphi, x)$ with $\bar{g}'(x)$ the solution of the Bayesian quickest disorder detection problem with linear delay penalty, under $\rho(x) \equiv \underline{\rho}$ and $\rho(x) \equiv \bar{\rho}$, respectively. In those cases, the problem of (2.16) degenerates into an optimal stopping problem for the one-dimensional Markov process $(\pi_t)_{t \geq 0}$ being equivalent to $(\varphi_t)_{t \geq 0}$, and the value functions $\underline{V}'(\pi, \varphi, x) \equiv \underline{V}'(\pi, \varphi) \equiv \underline{V}'(\varphi/(1 + \varphi), \varphi)$ and $\bar{V}'(\pi, \varphi, x) \equiv \bar{V}'(\pi, \varphi) \equiv \bar{V}'(\varphi/(1 + \varphi), \varphi)$ with the stopping boundaries $\underline{g}'(x) \equiv \underline{h}'$ and $\bar{g}'(x) \equiv \bar{h}'$ are given by the expressions in (4.15) and (4.14) below, whenever $\eta = 1/\underline{\rho}$ and $\eta = 1/\bar{\rho}$, respectively. Taking into account the properties of the boundary $g'(x)$ in (4.10) and the fact that $V'(\pi, \varphi, x) = 1 - \pi$ for all $\varphi \geq g'(x)$ and $0 \leq \pi < 1$, we therefore conclude by standard comparison arguments that the inequalities $\bar{V}'(\pi, \varphi) \leq V'(\pi, \varphi, x) \leq \underline{V}'(\pi, \varphi)$ and thus $0 < \lambda/c \leq \underline{h}' \leq g'(x) \leq \bar{h}'$ hold for all $(\pi, \varphi, x) \in [0, 1) \times [0, \infty) \times (0, \infty)$.

4.4. Let us finally introduce the function $\tilde{U}(\varphi/(1 + \varphi), \varphi, y) \equiv \tilde{G}(\varphi, y)$ and the boundary $\hat{h}(y)$ as a solution of the free-boundary problem consisting of the differential equation:

$$\left(\lambda(1 + \varphi) \frac{\partial G}{\partial \varphi} + \frac{\sigma^2(z e^{-\eta y} \varphi^\eta)}{\eta^2 z^2 e^{-2\eta y} \varphi^{2\eta}} \left(\frac{\varphi^2}{1 + \varphi} \frac{\partial G}{\partial \varphi} + \frac{\varphi^2}{2} \frac{\partial^2 G}{\partial \varphi^2} \right) \right) (\varphi, y) = -\frac{c\varphi}{1 + \varphi} \quad \text{for } \varphi < h(y) \quad (4.11)$$

instead of the one in (3.28), for each $y > 0$ fixed, and the conditions of (3.29)-(3.31) as well as (3.32)-(3.34) with $\phi = \varphi$ and $\pi = \varphi/(1 + \varphi)$. The general solution of the resulting first-order linear ordinary differential equation for $\varphi \mapsto (\partial G/\partial \varphi)(\varphi, y)$ takes the form:

$$\begin{aligned} \frac{\partial G}{\partial \varphi}(\varphi, y) &= \frac{C(y)}{(1 + \varphi)^2} \exp \left(\int_\varphi^w \frac{\lambda(1 + u)}{u^2} \frac{2\eta^2 z^2 e^{-2\eta y} u^{2\eta}}{\sigma^2(z e^{-\eta y} u^\eta)} du \right) \\ &\quad - \int_0^\varphi \frac{c(1 + u)}{u(1 + \varphi)^2} \frac{2\eta^2 z^2 e^{-2\eta y} u^{2\eta}}{\sigma^2(z e^{-\eta y} u^\eta)} \exp \left(- \int_u^\varphi \frac{\lambda(1 + v)}{v^2} \frac{2\eta^2 z^2 e^{-2\eta y} v^{2\eta}}{\sigma^2(z e^{-\eta y} v^\eta)} dv \right) du \end{aligned} \quad (4.12)$$

where $C(y)$ is an arbitrary continuously differentiable function, for each $y \in \mathbb{R}$ and any $z, w > 0$ fixed. By virtue of the assumptions of (2.2), we see that the term in the first line of (4.12)

above tends to infinity as $\varphi \downarrow 0$, so that $(\partial G/\partial \varphi)(\varphi, y) \rightarrow \pm\infty$ as $C(y) \neq 0$, for any $y \in \mathbb{R}$ fixed. We should thus choose $C(y) = 0$, that is equivalent to the property in (3.33). Hence, integrating the equation in (4.12), we therefore obtain that the solution of the system of (4.11) with (3.29) and (3.32)-(3.33) is given by:

$$G(\varphi, y; \tilde{h}(y)) = 1/(1 + \tilde{h}(y)) \quad (4.13)$$

$$+ \int_{\varphi}^{\tilde{h}(y)} \int_0^w \frac{c(1+u)}{u(1+w)^2} \frac{2\eta^2 z^2 e^{-2\eta y} u^{2\eta}}{\sigma^2(z e^{-\eta y} u^\eta)} \exp\left(-\int_u^w \frac{\lambda(1+v)}{v^2} \frac{2\eta^2 z^2 e^{-2\eta y} v^{2\eta}}{\sigma^2(z e^{-\eta y} v^\eta)} dv\right) du dw$$

for all $0 \leq \varphi < \tilde{h}(y)$, so that $0 \leq G(\varphi, y; \tilde{h}(y)) \equiv G(\varphi, y; z; \tilde{h}(y; z)) \leq 1/(1 + \varphi)$ holds, where $\tilde{h}(y)$ satisfies the equation:

$$\int_0^{\tilde{h}(y)} \frac{c(1+u)}{u} \frac{2\eta^2 z^2 e^{-2\eta y} u^{2\eta}}{\sigma^2(z e^{-\eta y} u^\eta)} \exp\left(-\int_u^{\tilde{h}(y)} \frac{\lambda(1+v)}{v^2} \frac{2\eta^2 z^2 e^{-2\eta y} v^{2\eta}}{\sigma^2(z e^{-\eta y} v^\eta)} dv\right) du = 1 \quad (4.14)$$

for each $y \in \mathbb{R}$ and any $z > 0$ fixed.

Summarizing these facts above, let us formulate the following assertion.

Corollary 4.4. *Suppose that $\mu_i(x)$, $i = 0, 1$, and $\sigma(x) > 0$ are continuously differentiable functions on $(0, \infty)$ in (2.1) satisfying (2.2) and (3.43)-(3.44) with $\alpha, \beta \in \mathbb{R}$ such that $(1 - \alpha)\eta \leq 0$ and $(1 - \beta)\eta \geq 0$, where $\eta = 1/(\eta_1 - \eta_0)$. Assume that $\tilde{h}(y)$ provides a unique solution of the equation in (4.14) for all $y \in \mathbb{R}$. Then, using the same arguments as in the proof of Lemma 3.3 above, it is shown that the function:*

$$\tilde{U}(\pi, \varphi, y) \equiv \tilde{G}(\varphi, y) = \begin{cases} G(\varphi, y; \tilde{h}(y)), & \text{if } 0 \leq \varphi < \tilde{h}(y) \\ 1/(1 + \varphi), & \text{if } \varphi \geq \tilde{h}(y) \end{cases} \quad (4.15)$$

coincides with the value function of the optimal stopping problem:

$$\tilde{U}(\pi, \varphi, y) = \inf_{\tau} E_{\pi, \varphi, y} \left[\frac{1}{1 + \varphi_\tau} + \int_0^\tau \left(\frac{c\varphi_t}{1 + \varphi_t} - \left(\frac{\lambda}{\varphi_t} + \lambda \right) \frac{\partial \tilde{G}}{\partial y}(\varphi_t, Y_t) I(\varphi_t < \tilde{h}(Y_t)) \right) dt \right] \quad (4.16)$$

with $\pi = \varphi/(1 + \varphi)$, which corresponds to the Bayesian risk function in (2.16). Moreover, $0 < \lambda/c \leq \tilde{h}(y) \equiv \tilde{h}(y; z)$ determined by (4.14) provides a hitting boundary for the stopping time:

$$\tilde{\tau} = \inf\{t \geq 0 \mid \varphi_t \geq \tilde{h}(Y_t)\} \quad (4.17)$$

which turns out to be optimal in (4.16) whenever the integral above is of finite expectation, for any $z > 0$ fixed.

Remark 4.5. Note that the function $\tilde{U}(\pi, \varphi, y)$ in (4.16) and the boundary $\tilde{h}(y)$ in (4.17) provide lower (upper) and upper (lower) estimates for the initial value function $U'(\pi, \varphi, y)$

defined as in (3.25) with $\alpha = 1$ and the optimal stopping boundary $h'(y)$ defined as in (3.26) with $\phi = \varphi$, whenever the function $y \mapsto \tilde{G}(\varphi, y)$ is increasing (decreasing) on \mathbb{R} . According to Remark 4.1 and the structure of the change of variables in (3.24) with $\phi = \varphi$, such a situation occurs when $\rho(x)$ from (3.12) is an increasing (decreasing) function on $(0, \infty)$ and $\eta_0 < \eta_1$ ($\eta_0 > \eta_1$) in (3.43), respectively.

Acknowledgments. This research was partially supported by Deutsche Forschungsgemeinschaft through the SFB 649 Economic Risk.

References

- [1] BAYRAKTAR, E. and DAYANIK, S. (2006). Poisson disorder problem with exponential penalty for delay. *Mathematics of Operations Research* **31**(2) (217–233).
- [2] BAYRAKTAR, E., DAYANIK, S. and KARATZAS, I. (2005). The standard Poisson disorder problem revisited. *Stochastic Processes and their Applications* **115**(9) (1437–1450).
- [3] BAYRAKTAR, E., DAYANIK, S. and KARATZAS, I. (2006). Adaptive Poisson disorder problem. *Annals of Applied Probability* **16** (1190–1261).
- [4] BEIBEL, M. (2000). A note on sequential detection with exponential penalty for the delay. *Annals of Statistics* **28** (1696–1701).
- [5] BENSOUSSAN, A. and LIONS, J. L. (1982). *Applications of Variational Inequalities in Stochastic Control*. (French Edition 1978) North Holland, Amsterdam.
- [6] DAYANIK, S. (2010). Compound Poisson disorder problems with nonlinear detection delay penalty cost functions. *Sequential Analysis* **29** (193–216).
- [7] DYNKIN, E. B. (1963). The optimum choice of the instant for stopping a Markov process. *Soviet Mathematical Doklady* **4** (627–629).
- [8] FRIEDMAN, A. (1976). *Stochastic differential equations and applications. Volumes I, II*. Academic Press, New York.
- [9] GAPEEV, P. V. and PESKIR, G. (2006). The Wiener disorder problem with finite horizon. *Stochastic Processes and their Applications* **116**(12) (1770–1791).

- [10] GAPEEV, P. V. and SHIRYAEV, A. N. (2010). On the sequential testing problem for some diffusion processes. To appear in *Stochastics: An International Journal of Probability and Stochastic Processes* (22 pp).
- [11] GRIGELIONIS, B. I. and SHIRYAEV, A. N. (1966). On Stefan's problem and optimal stopping rules for Markov processes. *Theory of Probability and its Applications* **11** (541–558).
- [12] KOLMOGOROV, A. N. (1992). On analitic methods in probability theory. *Selected works of A. N. Kolmogorov. Volume II. Probability Theory and Mathematical Statistics*. Shiryayev, A. N. ed. Kluwer, Dordrecht.
- [13] KRYLOV, N. V. (1980). *Controlled Diffusion Processes*. Springer, New York.
- [14] LIPTSER, R. S. and SHIRYAEV, A. N. (1977). *Statistics of Random Processes I*. Springer, Berlin.
- [15] ØKSENDAL, B. (1998). *Stochastic Differential Equations*. Springer, Berlin.
- [16] PESKIR, G. (2007). A change-of-variable formula with local time on surfaces. *Séminaire de Probabilité XL*. Lecture Notes in Mathematics **1899**, Springer (69–96).
- [17] PESKIR, G. and SHIRYAEV, A. N. (2006). *Optimal Stopping and Free-Boundary Problems*. Birkhäuser, Basel.
- [18] POOR, H. V. (1998). Quickest detection with exponential penalty for delay. *Annals of Statistics* **26** (2179–2205).
- [19] POOR, H. V. and HADJILIADIS, O. (2008). *Quickest Detection*. Cambridge University Press, Cambridge, United Kingdom.
- [20] POSPISIL, L., VECER, J. and HADJILIADIS, O. (2009). Formulas for stopped diffusion processes with stopping times based on drawdowns and drawups. *Stochastic Processes and thier Applications* **119** (2563–2578).
- [21] REVUZ, D. and YOR, M. (1999). *Continuous Martingales and Brownian Motion*. Springer, Berlin.
- [22] SHIRYAEV, A. N. (1961). The problem of the most rapid detection of a disturbance in a stationary process. *Soviet Mathematical Doklady* **2** (795–799).

- [23] SHIRYAEV, A. N. (1963). On optimum methods in quickest detection problems. *Theory of Probability and its Applications* **8** (22–46).
- [24] SHIRYAEV, A. N. (1964). On Markov sufficient statistics in nonadditive Bayes problems of sequential analysis. *Theory of Probability and its Applications* **9**(4) (670–686).
- [25] SHIRYAEV, A. N. (1965). Some exact formulas in a 'disorder' problem. *Theory of Probability and its Applications* **10** (348–354).
- [26] SHIRYAEV, A. N. (1967). Two problems of sequential analysis. *Cybernetics* **3** (63–69).
- [27] SHIRYAEV, A. N. (1978). *Optimal Stopping Rules*. Springer, Berlin.
- [28] SHIRYAEV, A. N. (2002). Quickest detection problems in the technical analysis of the financial data. *Mathematical Finance – Bachelier Congress 2000*. Selected papers from the First World Congress of the Bachelier Finance Society, Paris, June 29–July 1, 2000. Geman, H. et al. eds. Springer, Berlin (487–521).
- [29] SHIRYAEV, A. N. (2008). Generalized Bayesian nonlinear quickest detection problems: on Markov family of sufficient statistics. *Mathematical Control Theory and Finance*. Proceedings of the Workshop of April 10–14, 2007, Lisbon. Sarychev, A. et al. eds. Springer, Berlin (377–386).
- [30] SHIRYAEV, A. N. and ZRYUMOV, P. Y. (2009). On the linear and nonlinear generalized Bayesian disorder problem (discrete time case). *Optimality and Risk – Modern Trends in Mathematical Finance*. The Kabanov Festschrift. Delbaen, F. et al. eds. Springer, Berlin (227–235).
- [31] VERETENNIKOV, A. YU. (1980). On the strong solutions of stochastic differential equations. *Theory of Probability and its Applications* **24** (354–366).